



# **MANUAL OF THE FRESHWATER BIVALVES OF MARYLAND**



**CHESAPEAKE BAY AND  
WATERSHED PROGRAMS**  
MONITORING AND  
NON-TIDAL ASSESSMENT  
CBWP-MANTA- EA-96-03





# **MANUAL OF THE FRESHWATER BIVALVES OF MARYLAND**

Prepared By:

Art Bogan<sup>1</sup> and Thomas Proch<sup>2</sup>

<sup>1</sup>Freshwater Molluscan Research  
36 Venus Way  
Sewell New, Jersey 08080-1970

<sup>2</sup>Pennsylvania Department of Environmental Protection  
400 Waterfront Drive  
Pittsburgh, PA 15222-4745

Prepared For:

John Christmas  
Maryland Department of Natural Resources  
Chesapeake Bay and Watershed Programs  
Resource Assessment Service  
Monitoring and Non-Tidal Assessment Division  
Ecological Assessment Program  
580 Taylor Avenue  
Annapolis, Maryland 21401

January 1997





# Table of Contents

I. Schedule of Workshop	1
II. List of maps	2
III. List of figures	2
IV. Introduction	3
V. Acknowledgments	4
VI. Figure of bivalve shell landmarks (fig. 1)	5
VII. Glossary of bivalve terms	6
VIII. Drainage basin map of Maryland (map 1)	9
IX. List of freshwater bivalve families occurring in Maryland	10
X. List of unionids of Maryland with common and scientific names and status	11
XI. List of Sphaeriidae of Maryland with common and scientific names and status	11
XII. Key to the freshwater bivalves of Maryland	12
XIII. List of Institutions with major unionid bivalve collections	15
XIV. Species descriptions:	
1. <i>Alasmidonta heterodon</i> (Lea, 1829)	17
2. <i>Alasmidonta marginata susquehannae</i> Ortmann, 1919	19
3. <i>Alasmidonta undulata</i> (Say, 1817)	21
4. <i>Alasmidonta varicosa</i> (Lamarck, 1819)	23
5. <i>Anodonta implicata</i> Say, 1829	25
6. <i>Elliptio angustata</i> (Lea, 1831)	27
7. <i>Elliptio complanata</i> (Lightfoot, 1786)	29
8. <i>Elliptio fisheriana</i> (Lea, 1838)	33
9. <i>Lampsilis cardium</i> Rafinesque, 1820	35
10. <i>Lampsilis cariosa</i> (Say, 1817)	37
11. <i>Lampsilis radiata</i> (Gmelin, 1791)	39
12. <i>Lasmigona subviridis</i> (Conrad, 1835)	41
13. <i>Leptodea ochracea</i> (Say, 1817)	43
14. <i>Ligumia nasuta</i> (Say, 1817)	45
15. <i>Pyganodon</i> (= <i>Anodonta</i> ) <i>cataracta</i> (Say, 1817)	47
16. <i>Strophitus undulatus</i> (Say, 1817)	49
17. <i>Utterbackia</i> (= <i>Anodonta</i> ) <i>imbecillis</i> (Say, 1829)	51
XV. Color plates of freshwater bivalves (figs. 2-20)	53

**XVI. Introduction to the literature on freshwater bivalves**

1. Sphaeriidae . . . . .	56
2. Corbiculidae . . . . .	56
3. Dreissenidae . . . . .	56
4. Unionoidea. . . . .	58
a. Handbook, field guides and regional faunas . . . . .	58
b. Taxonomy . . . . .	60
c. Literature on Maryland unionids . . . . .	61
d. Endangered species . . . . .	62
e. Glochidia, reproduction and fish hosts . . . . .	64
f. Food . . . . .	67

XVII. Appendix 1. Freshwater bivalves reported from the Youghiogheny and Casselman rivers in Pennsylvania and potentially occurring in Maryland . . . . .	68
---	----

## List of Maps

Map	1	Drainage Basins of Maryland . . . . .	9
Map	2	<i>Alasmidonta heterodon</i> (Lea, 1829) . . . . .	16
Map	3	<i>Alasmidonta marginata susquehannae</i> Ortmann, 1919 . . . . .	18
Map	4	<i>Alasmidonta undulata</i> (Say, 1817) . . . . .	20
Map	5	<i>Alasmidonta varicosa</i> (Lamarck, 1819) . . . . .	22
Map	6	<i>Anodonta implicata</i> Say, 1829 . . . . .	24
Map	7	<i>Elliptio angustata</i> (Lea, 1831) . . . . .	26
Map	8	<i>Elliptio complanata</i> (Lightfoot, 1786) . . . . .	28
Map	9	<i>Elliptio fisheriana</i> (Lea, 1838) . . . . .	32
Map	10	<i>Lampsilis cardium</i> Rafinesque, 1820 . . . . .	34
Map	10	<i>Lampsilis cariosa</i> (Say, 1817) . . . . .	36
Map	12	<i>Lampsilis radiata</i> (Gmelin, 1791) . . . . .	38
Map	13	<i>Lasmigona subviridis</i> (Conrad, 1835) . . . . .	40
Map	14	<i>Leptodea ochracea</i> (Say, 1817) . . . . .	42
Map	15	<i>Ligumia nasuta</i> (Say, 1817) . . . . .	44
Map	16	<i>Pyganodon</i> (= <i>Anodonta</i> ) <i>cataracta</i> (Say, 1817) . . . . .	46
Map	17	<i>Strophitus undulatus</i> (Say, 1817) . . . . .	48
Map	18	<i>Utterbackia</i> (= <i>Anodonta</i> ) <i>imbecillis</i> (Say, 1829) . . . . .	50

## List of Figures

Figure	1	Bivalve shell landmarks . . . . .	5
Figure	2	<i>Dreissena polymorpha</i> (Pallas, 1771) . . . . .	53
Figure	3	<i>Corbicula fluminea</i> (Müller, 1776) . . . . .	53
Figure	4	<i>Alasmidonta heterodon</i> (Lea, 1829) . . . . .	53
Figure	5	<i>Alasmidonta marginata susquehannae</i> Ortmann, 1919 . . . . .	53
Figure	6	<i>Alasmidonta undulata</i> (Say, 1817) . . . . .	53
Figure	7	<i>Alasmidonta varicosa</i> (Lamarck, 1819) . . . . .	53
Figure	8	<i>Anodonta implicata</i> Say, 1829 . . . . .	54
Figure	9	<i>Elliptio angustata</i> (Lea, 1831) . . . . .	54
Figure	10	<i>Elliptio complanata</i> (Lightfoot, 1786) . . . . .	54
Figure	11	<i>Elliptio fisheriana</i> (Lea, 1838) . . . . .	54
Figure	12	<i>Lampsilis cardium</i> Rafinesque, 1820 . . . . .	54
Figure	13	<i>Lampsilis cariosa</i> (Say, 1817) . . . . .	54
Figure	14	<i>Lampsilis radiata</i> (Gmelin, 1791) . . . . .	54
Figure	15	<i>Lasmigona subviridis</i> (Conrad, 1835) . . . . .	55
Figure	16	<i>Leptodea ochracea</i> (Say, 1817) . . . . .	55
Figure	17	<i>Ligumia nasuta</i> (Say, 1817) . . . . .	55
Figure	18	<i>Pyganodon</i> (= <i>Anodonta</i> ) <i>cataracta</i> (Say, 1817) . . . . .	55
Figure	19	<i>Strophitus undulatus</i> (Say, 1817) . . . . .	55
Figure	20	<i>Utterbackia</i> (= <i>Anodonta</i> ) <i>imbecillis</i> (Say, 1829) . . . . .	55



## INTRODUCTION

The freshwater bivalves of the state of Maryland are represented by 1 introduced species and historically by 17 native unionid species in the Chesapeake Bay Basin and by 22 species prehistorically in the Monongahela River Basin. The only comprehensive monograph of the freshwater bivalves of the region of including Maryland and Pennsylvania was prepared by Arnold E. Ortmann (1913, 1919), while working at The Carnegie Museum of Natural History, Pittsburgh. This workbook is designed to provide an introduction to the freshwater bivalves of Maryland. This is accomplished by providing a list of all of the freshwater bivalves, excluding Sphaeriidae, their common names and distribution by river basin. A key based on shell characters for all of the species, except Sphaeriidae, is provided. Each unionid species has a detailed shell description and is supplemented by a color figure of a representative specimen of the species. A glossary is included to obviate any obscure descriptive terms used in the notebook. An introductory bibliography is included to facilitate entry into the literature on all three families of freshwater bivalves occurring in the state. The bibliography is broken down by family and is subdivided by major topic under Unionoidea.

The purpose of the workshop is to train regulatory agency personnel and other interested parties in recognition and identification of freshwater bivalves that either are known or suspected to occur in Maryland. It is hoped that such training will lead to increased knowledge of species distribution in the state. During the workshop participants will be exposed to the diversity of North American freshwater bivalves and the importance of these animals in aquatic ecology. There will be sufficient material of each species to examine so that participants will have an opportunity to begin to develop an idea of the range of variation within species reported from the state. The combined use of the key, figures and descriptions of the species should help the participants to learn which characters are important in the identification of shells of these species.

The format for each of the species accounts includes: a map of the historic distribution of the species but does not include the most recent survey data (eg. Bartgis and MacIvor, 1993), the basic map was modified from Gerberich (1984); common names follow Turgeon et al. (1988); synonymy includes junior synonyms and previous generic combinations; shell description is a detailed list of the characteristics of the shell of the species described; distribution is the list of states in which the species occurred historically based on Williams et al. (1993); ecology is the published notes on the ecology of the species; breeding season is that period of the year when female specimens have been observed with glochidia in the marsupium; fish hosts are noted where known as compiled from the literature by Watters (1994); status is the perceived status of the species throughout its range as presented by Williams et al. (1993).

### **ACKNOWLEDGMENTS**

The bibliographic citations used in this volume were taken from the author's work in progress on the bibliography of unionids of North America with Kevin Cummings and Clement L. Counts, III. The use of this bibliography greatly facilitated the development of the literature section. John Christmas was instrumental in developing the workshop and revising the key to the freshwater bivalves of Maryland. Dr. John Rawlins, Curator of Invertebrates, Carnegie Museum of Natural History, Pittsburgh provided access to the collection for the photography of several species of unionids and provided specimens for the workshop. The Delaware Museum of Natural History and Dr. Paula M. Mikkelsen, Curator of Malacology graciously provided study specimens used during the workshop.




Figure 1. Morphology of a freshwater mussels shell illustrating structures and terminology. a. exterior of right valve, b. interior of left valve. (from Burch 1975:5, fig.2; Bogan and Parmalee 1983:9).



## GLOSSARY OF BIVALVE TERMS

- Alate - with an extension or wing on the dorsal edge of the shell.
- Angular (subangulate) - having either the anterior or posterior margins forming a relatively acute (sharp) angle.
- Anterior - front or forward.
- Arcuate - bent in a bow or arched.
- Beak - the raised portion of the dorsal margin of a shell; formed by the embryonic shell around which the rest of the shell develops distally in a concentric manner.
- Beak cavity - the cavity on the inside of each valve leading into the beak, under the interdentum.
- Beak sculpture - raised ridges or undulations on the umbo.
- Biangular - having two angles.
- Bradytic - mussels which are long term breeders; females retain glochidia in their gills typically over the winter.
- Byssus, byssal threads - a bundle of tough threads secreted by the byssal gland in the foot of a bivalve, used to anchor the bivalve to some hard substrate.
- Cardinal teeth - teeth located between the lateral teeth in Corbiculidae and Sphaeriidae.
- Chevron - shaped like a wide-angled V.
- Clinal variation - the graded variation in morphology exhibited by a species in mollusks from headwater areas to the mouth of the highest order stream.
- Compressed (subcompressed) - flattened out or pressed together.
- Concentric - having a common center, such as ridges or loops radiating from the beak of a mussel valve.
- Conspecific - pertaining to individuals or populations of the same species.
- Corrugated - marked by wrinkles or ridges and grooves.
- Crescentic - shaped like the figure of the crescent moon with a convex and a concave edge.
- Decorticate - to remove the outer covering, in mollusks the epidermis.
- Disc - the middle or central portion of the exterior of a valve; distinct from the posterior slope or other areas immediately adjacent to the margin of the valve.
- Discoidal - round and flat like a disc.
- Dorsal - the top or back; in mussels, the hinge area.
- Edentulous - lacking both pseudocardinal and lateral teeth.
- Effuse - spread out broadly.
- Elliptical (subelliptical) - elongated, having the form of an ellipse.
- Elongate - long or extended.
- Emarginate - having a shallow notching at the margin.
- Endangered - this status at the state level includes peripheral forms which may be common in another part of its range, but whose continued existence within the political boundaries of the state is in danger of extirpation. At the national level, this status means the organism is in danger of extinction, and included on or being considered for the U.S. List of Endangered Fauna and Endangered and Threatened Plant Species of the United States, under the Endangered Species Act of 1973 (Cooper et al. 1973:x).
- Epidermis - exterior or outside (corneous) layer of the shell.
- Extinct - a species which has no living representatives; all individuals are no longer extant.
- Extirpated - the extinction of a species within a portion of its range.
- Form - an animal with questionable taxonomic status; that is, one exhibiting variation but the extent or degree is not well enough known to determine whether it is a species, subspecies or simply individual or population variation.
- Fusiform - tapering toward each end.
- Gills - a thin plate-like paired structure within the mantle cavity which serves as a respiratory organ in aquatic mollusks and in female unionids all of the gills or certain portions of the gills serve as the marsupium.
- Globose - globe-like, spherical.
- Glochidium (plural - glochidia) - the bivalve larvae of unionids which are generally parasitic on the gills of fish.
- Gravid female - a female which has embryos in the marsupium.
- Growth lines - compact lines of temporarily arrested growth or rest periods appearing on the epidermis of the shell as a raised or darker concentric line.
- Hinge ligament - an elastic, elongate, corneous structure that unites the two valves dorsally along the hinge plate.
- Holotype - single specimen designated as the "type" by the author in the publication of a new species level taxon.



- Inequilateral - in a bivalve, having the two ends unequal, i.e., one end is wider or thicker than the other.
- Inflated (subinflated) - moderately to greatly inflated.
- Interdentum - a flattened area of the hinge plate between the pseudocardinal and lateral teeth.
- Iridescent - showing colors like those of a rainbow.
- Lachrymose - term describing teardrop-shaped pustules.
- Lateral teeth - the elongated, raised and interlocking structures along the hinge line of the valve.
- Lectotype - one of a series of syntypes which, subsequent to the publication of an original description of a species level taxon, is selected (by publication) to serve as the type specimen for that taxon.
- Lunule - depressed area immediately anterior to the umbo.
- Marsupial swelling - a section of the posterior ventral margin of the female unionid shell which is enlarged or inflated to provide space for expansion of the marsupium with the development of the glochidia.
- Marsupium (marsupial pouch) - in unionids, a brood pouch for eggs and developing glochidia, formed by a restricted portion of the outer gill, the complete outer gill or all four gills.
- Muscle scar - the area of attachment of a muscle to the inside of the shell; e.g., the anterior adductor muscle scar is the location of attachment for the anterior adductor muscle.
- Nacre - the interior iridescent, thin layer of a mussel shell.
- Naiad - formerly a tribe of mollusca nearly equivalent taxonomically to the family Unionidae, often used as a synonym of unionid.
- Nodule (subnodulous) - a small rounded mass of irregular shape.
- Oblique - slanting; angled, but not horizontal or vertical.
- Obovate - (subobovate) - ovate.
- Orbicular (suborbicular) - having the form of an orb; circular or nearly circular in outline.
- Oval, Ovate (subovate) - egg-shaped, broadly elliptical.
- Pallial line - an indented groove or line approximately parallel with the ventral margin of a bivalve shell which marks the line of muscles attaching the mantle to the shell.
- Paratype - each specimen of a type series other than the holotype designated in the original publication of the taxon.
- Periphery - the external boundary on a surface, edge.
- Periostracum - see epidermis.
- Plications - parallel ridges on the surface of the shell.
- Posterior - hind or rear.
- Posterior ridge - a ridge on the exterior of a mussel shell, extending from the umbo to the posterior margin.
- Posterior slope - the area across the dorsal portion of the valve extending from the umbo to the posterior margin, often above the posterior ridge.
- Pseudocardinal teeth - triangular-shaped hinge teeth near the anterior -dorsal margin of the shell.
- Pustule - small, raised structure on the external or outside surface of the shell (see also tubercle).
- Quadrangle (subquadrangle) - square, or nearly square in outline.
- Radial furrow - a groove or depression; in naiads a groove running from the umbo area toward the shell margin.
- Radiating - proceeding outward from a central point.
- Rare - seldom appearing, occurring widely separated in space; extremely few in number.
- Rectangular - a shape with four sides possessing four right-angles.
- Rest mark - see growth lines.
- Rhomboid (subrhomboid) - having generally four distinct sides, two sides being longer than the others.
- Semicircular - a partial or incomplete circle.
- Serrated - notched or grooved.
- Sexual dimorphism - a condition in which males and females of the same species are morphologically different, usually indicated by an expanded posterior marsupial area in the female in contrast to a more pointed or bluntly rounded area in the male.
- Sinus - a character of some unionids which have a depression above or below the posterior ridge.
- Solid (subsolid) - shells which are thick and heavy.
- Special Concern - This status covers cases where the organism exists in small populations over a broad range, may be over exploited which may pose a threat, the organism are especially vulnerable to specific pressures, or any other reasons identified by experienced researchers (Cooper et al. 1973:x).
- Species - group of interbreeding natural populations that are reproductively isolated from all other such groups.
- Striae - impressed or raised lines on a shell.

Striate - having striae.

Subspecies - a geographically defined aggregate of local populations within a species which differ morphologically and/or physiologically from other aggregations of local populations within that species.

Sulcus (plural - sulci) - a longitudinal furrow or depression.

Sympatric - pertaining to populations of two or more closely related species which occupy identical or broadly overlapping geographical areas.

Syntype - one of a series of specimens of the same taxon which formed the material studied by the original author to describe a new species level taxon, from which no type specimen (holotype) was designated.

Tachytictic - mussels which are short-term breeders; i.e., glochidia are found in the gills of the female only during the summer.

Taxon - any formal taxonomic unit or category of an organism; e.g., a species or genus.

Threatened - This status at the state level includes forms which are likely to become Endangered in the foreseeable future if certain conditions are not met. This includes forms which exhibit a considerable decrease in numbers beyond normal populations fluctuations or a documented range contraction, but are not yet considered Endangered. At the national level this applies to the Endangered Species Act of 1973 (Cooper et al. 1973:x).

Trapezoid (subtrapezoid) - a shape having four distinct sides with two sides parallel.

Triangular (subtriangular) - a shape having three sides and three angles, like a triangle.

Truncate (subtruncate) - having the end squared off.

Tubercle (tuberculate) - small, raised, rounded knob on the outside of the shell.

Tuberculate - having tubercles on the outside of the shell.

Type - a designated specimen or specimens of an organism that serves as the basis for the original name and description of any species level taxon.

Umbo/umbone - the dorsally raised, inflated area of the bivalve shell.

Unionids - refers to any member of the freshwater bivalve mollusks which belong to the superfamily Unionoidea.

Undulation - pattern with waves; raised ridges or bars.

Valve - the right or left half of a mussel (or unionid) shell.


Ventral - the underside or bottom.

The following volumes, in addition to *Webster's Unabridged Dictionary*, were used to compile the definitions used in the glossary.

- Burch, John B. 1962. *How to know the eastern land snails*, Pictured Key Nature Series, Wm. C. Brown Company Publishers, Dubuque, Iowa, 214 pp. 519 fig.
- Burch, John B. 1975. *Freshwater Unionacean Clams (Mollusca: Pelecypoda) of North America*. Malacological Publications, Hamburg, Michigan, 204 pp. 252 fig.
- Leonard, A. Byron. 1954. Handbook of Gastropods in Kansas. *University of Kansas Museum of Natural History Miscellaneous Publication No. 20*, pp. 1-224, 22 pl.
- Murray, Harold D. and A. Byron Leonard. 1962. Handbook of Unionid Mussels in Kansas, *University of Kansas Museum of Natural History Miscellaneous Publication No. 28*, pp. 1-184, 45 pl. 42 fig.
- Parmalee, Paul W. 1967. The Freshwater Mussels of Illinois. *Illinois State Museum, Popular Science Series*, Vol. 8, pp 108, 35 pl. 4 fig.
- Pennak, Robert W. 1964. *Collegiate dictionary of zoology*, The Ronald Press Company, New York. 583 pp.

This glossary was taken from Bogan (1993).

Map 1. Eighteen Major Drainage basins of Maryland (adapted from printout from Maryland Department of Natural Resources)



**List of families/superfamilies of freshwater bivalves occurring Maryland including the higher molluscan classification.**

Phylum Mollusca

Class Bivalvia

Subclass Paleoheterodonta

Order Unionoida

Superfamily Unionoidea

Family Unionidae [10 genera, 17 species] (see page 11 for species list)

<i>Alasmidonta</i>	[4 species]
<i>Anodonta</i>	[1]
<i>Elliptio</i>	[3]
<i>Lampsilis</i>	[3]
<i>Lasmigona</i>	[1]
<i>Leptodea</i>	[1]
<i>Ligumia</i>	[1]
<i>Pyganodon</i>	[1]
<i>Strophitus</i>	[1]
<i>Utterbackia</i>	[1]

Subclass Heterodonta

Order Veneroida

Superfamily Dreissenoidea

Family Dreissenidae

*Dreissena* [Zebra mussel, quagga mussel] **Not yet known from Maryland!**

Superfamily Corbiculoidea

Family Corbiculidae

*Corbicula fluminea* (Müller, 1774) [Asian clam]

Family Sphaeriidae [fingernail clams or pillclams] (3 genera, 20 species; see page 11)

<i>Musculium</i>	[4]
<i>Pisidium</i>	[12]
<i>Sphaerium</i>	[4]

## List of unionid bivalves of Maryland with common and scientific names and status

<u>Species</u>	<u>Common Name</u> <sup>1</sup>	<u>Status</u>			
		*	X	E	S
<i>Alasmidonta heterodon</i> (Lea, 1829)	dwarf wedgemussel	*		E	
<i>Alasmidonta marginata susquehannae</i> Ortmann, 1919	Susquehanna elktoe		?		
<i>Alasmidonta undulata</i> (Say, 1817)	triangle floater			E	
<i>Alasmidonta varicosa</i> (Lamarck, 1819)	brook floater		X	E	
<i>Anodonta implicata</i> Say, 1829	alewife floater				
<i>Elliptio angustata</i> (Lea, 1831)	Carolina lance			E	
<i>Elliptio complanata</i> (Lightfoot, 1786)	eastern elliptio				
<i>Elliptio fisheriana</i> (Lea, 1838)	northern lance				S
<i>Lampsilis cardium</i> Rafinesque, 1820	plain pocketbook				
<i>Lampsilis cariosa</i> (Say, 1817)	yellow lampmussel			E	
<i>Lampsilis radiata</i> (Gmelin, 1791)	eastern lampmussel				
<i>Lasmigona subviridis</i> (Conrad, 1835)	green floater			E	
<i>Leptodea ochracea</i> (Say, 1817)	tidewater mucket				
<i>Ligumia nasuta</i> (Say, 1817)	eastern pondmussel				
<i>Pyganodon (=Anodonta) cataracta</i> (Say, 1817)	eastern floater				
<i>Strophitus undulatus</i> (Say, 1817)	squawfoot				
<i>Utterbackia (=Anodonta) imbecillis</i> (Say, 1829)	paper pondshell				

## List of sphaeriidae of Maryland with common and scientific names (Gerberich, 1984).

<u>Species</u>	<u>Common Name</u>	<u>Status</u>			
		*	X	E	S
<i>Musculium lacustre</i> (Müller, 1774)	lake fingernailclam				
<i>Musculium partumeium</i> (Say, 1822)	swamp fingernailclam				
<i>Musculium securis</i> (Prime, 1852)	pond fingernailclam				
<i>Musculium transversum</i> (Say, 1829)	long fingernailclam				
<i>Pisidium adamsi</i> Stimpson, 1851	Adam peaclam				
<i>Pisidium casertanum</i> (Poli, 1791)	ubiquitous peaclam				
<i>Pisidium compressum</i> Prime, 1852	ridged-beak peaclam				
<i>Pisidium dubium</i> (Say, 1817)	greater eastern peaclam				
<i>Pisidium equilaterale</i> Prime, 1852	round peaclam				S
<i>Pisidium fallax</i> Sterki, 1896	river peaclam				
<i>Pisidium ferrugineum</i> Prime, 1852	rusty peaclam				
<i>Pisidium lilljeborgi</i> (Clessin, 1886)	Lilljeborg peaclam				
<i>Pisidium nitidum</i> Jenyns, 1852	shiny peaclam				
<i>Pisidium punctatum</i> Sterki, 1895	perforated peaclam				
<i>Pisidium variable</i> Prime, 1852	triangular peaclam				
<i>Pisidium walkeri</i> Sterki, 1895	Walker peaclam				
<i>Sphaerium fabale</i> (Prime, 1852)	river fingernail clam				
<i>Sphaerium occidentale</i> (Lewis, 1856)	Herrington fingernailclam				
<i>Sphaerium simile</i> (Say, 1817)	grooved fingernailclam				
<i>Sphaerium striatinum</i> (Lamarck, 1818)	striated fingernailclam				

\* federally endangered

X presumed extinct in Maryland

E suggested as endangered in Maryland

S suggested as special concern in Maryland

<sup>1</sup> Common names follow Turgeon et al. (1988)

*Freshwater Bivalves of Maryland*

**KEY TO THE SHELLS OF THE FRESHWATER BIVALVES OF MARYLAND (1/97 update)**

- 1
  - a. Shell with very sharp posterior ridge, shaped like a marine mussel, *Mytilus*, generally less than 30 millimeters, and attached to a hard substrate with byssal threads (fig.2)  
.....*Dreissena polymorpha*  
*Note: (nonindigenous aquatic nuisance species, not-found in wild in Maryland)*
  - b. Animal without byssal threads attaching adult animal to substrate, with or without teeth but not with above shape.....2
- 2 (1b)
  - a. Valves with cardinal teeth and two sets of lateral teeth.....3
  - b. Valves with one set of lateral teeth and pseudocardinal teeth or without teeth (Unionidae).....4
- 3 (2a)
  - a. Valves with serrated lateral teeth (fig. 3), nonindigenous species.....*Corbicula fluminea*
  - b. Valves with smooth lateral teeth.....*Sphaeriidae*
- 4 (2b)
  - a. Specimens from the Ohio basin.....Appendix A
  - b. Specimens from the Chesapeake or coastal basins.....5
- 5 (4b)
  - a. Hinge teeth absent.....6
  - b. Hinge teeth present.....9
- 6 (5a)
  - a. Beaks not projecting above the hinge line (fig.20).....*Utterbackia imbecillis*
  - b. Beaks projecting above the hinge line.....7  
1 (footnote)
- 7 (6b)
  - a. Beak sculpture double looped<sup>1</sup>.....8
  - b. Vestige of pseudocardinal teeth usually represented by a thickening near the beaks (fig.19); nacre usually orange in the beak cavity, beak sculpture concentric  
.....*Strophitus undulatus*
8. (7b)
  - a. Shell prominently thickened along the anterior third of the ventral margin of the shell below the pallial line, nacre salmon or copper colored (fig. 8) .....*Anodonta implicata*
  - b. Ventral margin of shell uniformly thin, nacre bluish or white, greenish epidermis (fig.18).....*Pyganodon cataracta*
- 9 (5b)
  - a. Lateral teeth well developed, functional and interlocking.....12
  - b. Lateral teeth absent or reduced, neither functional nor interlocking.....10
- 10 (9b)
  - a. Fine transverse ridges present on the posterior slope, pseudocardinal teeth reduced and elongate, with smooth surfaces.....11
  - b. Transverse ridges on posterior slope absent, pseudocardinal teeth strong and triangular, with rough surfaces, shell small to medium size, triangular to ovate (fig.6).....



# *Freshwater Bivalves of Maryland*

- .....*Alasmidonta undulata*
- 11(10a) a. Posterior ridge angular and prominent, shell truncated, color of posterior margin lighter than rest of shell (fig.5).....*Alasmidonta marginata susquehannae*
- b. Posterior ridge rounded, shell rounded and usually less than 70 mm long (fig.7)  
          .....*Alasmidonta varicosa*
- 12 (9a) a. Right valve with two lateral teeth, rare (fig.4).....*Alasmidonta heterodon*
- b. Right valve with one lateral tooth.....13
- 13 (12b) a. Height/length ratio less than or equal to 0.5 shell elongate.....14
- b. Height/length ratio greater than 0.5.....15
- 14 (13a) a. Posterior ridge prominent, posterior end of shell subangular; pseudocardinal teeth thin, blade-like, some shells with prominent rays, shell usually less than 110 mm in length (fig. 17).....*Ligunia nasuta*
- b. Dark brown periostracum posterior ridge rounded, posterior end bluntly rounded, ventral margin broadly rounded lacks rays (fig.11), most commonly distributed lanceolate species on Maryland's Eastern Shore.....*Elliptio fisheriana*
- c. Posterior ridge rounded, posterior end bluntly rounded, ventral margin broadly rounded lacks rays (fig.11), most commonly distributed lanceolate species in Potomac drainage and Central part of state. ....*Elliptio producta*
- d. Posterior ridge rounded, posterior end bluntly rounded, ventral margin broadly rounded lacks rays (fig.11), periostracum waxy, ranging from green to yellow.....*Elliptio lanceolata*
- e. Shell elongate, with a well-defined posterior ridge, posterior end bluntly pointed, ventral margin straight (fig. 9).....*Elliptio angustata*
- 15 (13b) a. Nacre purple, may grade to a salmon color, shell subrhomboid with well-defined posterior ridge and slope, very common (fig.10) periostracum yellowish green to black.....*Elliptio complanata*
- b. Nacre white or colored but not purple.....16
- 16(15b) a. Left valve with small interdental tooth, giving the appearance of three pseudocardinal teeth, shell more or less compressed and subrhomboid in outline, periostracum dark green with numerous green rays or brown, adult shell less than 65 mm long, posterior ridge rounded, beak sculpture is prominent raised bars (fig.15).....*Lasmygona subviridis*
- b. Left valve without interdental tooth, valve appears to have only two pseudocardinal teeth.....17
- 17(16b) a. Adult shell usually less than 80 mm in length, and thin, hardly thicker anteriorly than posteriorly, periostracum dull yellow without rays or with fine rays all over the shell, in or near tidewaters, nacre often a salmon color (fig.16).....*Leptodea ochracea*

# Freshwater Bivalves of Maryland

- b. Adult shell often greater than 80 mm in length, much thicker anteriorly than posteriorly, may have obvious broad color rays.....19
- 18 (17b) a. Over 1.5 times as long as high. Height/length less than 0.60 mm or cm in males and in most females, shell with rays all over the shell (may be obscured in old adults), posterior ridge low and broadly rounded, beaks not prominent (fig. 14).....*Lampsilis radiata*
- b. Height/length greater than 0.60, mm or cm beaks prominent, periostracum yellow to light brown now bluish white.....20
- 19 (18b) a. Periostracum glossy yellow reddish brown in adults. Green rays if present are very thin and restricted to the posterior slope, posterior ridge very smooth and rounded pseudocardinal teeth are both serrated and perpendicular to the hinge line, pseudocardinal teeth on left valve are perpendicular to hinge line. (fig.13).....*Lampsilis cariosa*
- b. Periostracum yellow to yellowish-brown, sometimes marked with distinct dark green rays, which can be thicker than those in *L. cariosa* (indistinct in old shells), shell globose, almost as high as long, posterior ridge rounded, beaks very inflated (fig.12) nacre silver white and pink under beak, pseudocardinal teeth on left valve are parallel to hinge line, nonindigenous species, introduced into the Potomac River..... *Lampsilis cardium*<sup>3</sup>

<sup>1</sup>Beak sculpture is often difficult to evaluate, as a result of shelf erosion, in all but very small specimens.

<sup>2</sup>There are taxonomic problems with the group of lanceolate species in the genus *Elliptio*. Several names have been used for this group on Maryland including *E. fisheriana* (Lea 1838), *E. producta* (Conrad, 1836) and *E. angustata* (Lea, 1831). The last two species were described from the Savannah River in South Carolina and Georgia and the Cooper and Congaree rivers in South Carolina, respectively. *E. fisheriana* (Lea, 1838) was described from the "head of Chester River, [Kent Co.] Maryland." There is confusion as to what constitutes a species and where one ends and the next begins. *E. angustata* was described first so probably should be the name used for this shell shape until this species complex is resolved. *E. producta* has also been described in Maryland.

<sup>3</sup>The inclusion of *Lampsilis ovata* (Say, 1817) on the list of species for Maryland is in error. It stems from the placement of the subspecies Ortmann described, *ventricosa cohongoronta* Ortmann, 1912, from the Potomac River. The confusion is a result of some people using *ventricosa* (Barnes, 1823) as either a subspecies or junior synonym of *L. ovata*, it is not. *Cardium* Rafinesque, 1820 is an earlier name and senior synonym for *ventricosa* (Barnes, 1823).



**List of Institutions with major unionid collections, address and contact person.**

Dr. John B. Burch  
Malacology  
Museum of Zoology  
University of Michigan  
Ann Arbor, Michigan 48109

Dr. Robert Hershler  
Division of Mollusks  
United State National Museum  
Smithsonian Institution  
Washington, DC 20560

Dr. Ken Boss  
Division of Mollusks  
Museum of Comparative Zoology  
Harvard University  
Cambridge, MA 02138

Dr. Paula M. Mikkleson  
Mollusks  
Delaware Museum of Natural History  
P.O. Box 4937  
Wilmington, DE 19807-0937


Dr. Paul W. Parmalee  
McClung Museum  
University of Tennessee  
Knoxville, TN 37916

Dr. John E. Rawlins  
Division of Invertebrates  
Carnegie Museum of Natural History  
4400 Forbes Ave.  
Pittsburgh, PA 15213

Dr. Gary Rosenberg  
Department of Malacology  
Academy of Natural Sciences  
1900 Benajamin Franklin Parkway  
Philadelphia, PA 19103

Dr. David H. Stansbery  
Museum of Biological Diversity  
1315 Kinnear Road  
Columbus, OH 43212-1192

Dr. Fred Thompson  
Mollusks  
Florida Museum of Natural History  
University of Florida  
Gainesville, FL 32611



Map 2      Distribution of *Alasmidonta heterodon* (Lea, 1829) in Maryland.

***Alasmidonta heterodon* (Lea, 1829) dwarf wedgemussel Fig. 4****SYNONYMY.**

*Unio heterodon* Lea, 1829

*Alasmidonta heterodon* (Lea, 1829)

*Alasmidonta (Pressodonta) heterodon* (Lea, 1829)

**SHELL DESCRIPTION.** Shell small, subtrapezoid or "hump backed" shell thick anteriorly and thinning posteriorly, ventral margin mostly straight, posterior margin pointed near the base, dorsal margin slightly curved, beaks low and rounded, projecting only slightly above the hinge line, posterior ridge rounded, somewhat inflated and prominent, periostracum yellowish, olive brown to blackish with variable width reddish brown rays, hinge teeth unusual with an interdental projection, beak cavity narrow and rather shallow, nacre bluish white.

**DISTRIBUTION.** Connecticut, Delaware, Massachusetts, Maryland, North Carolina, New Hampshire, New Jersey, New York, Pennsylvania, Virginia, Vermont, Canada: New Brunswick (Williams et al. 1993).


**ECOLOGY.** Ortmann (1919:176) observed "I found this species in very small streams ("runs"), in strongly flowing water, and in rather coarse gravel, ... in a small branch of the river in sand and moderate current ... in the canal at Manayunk at time when the water had been drained off ... Here the bottom consisted of larger and smaller stones, the interstices filled with sandy mud. This however, cannot be regarded as normal, and the ecological conditions of this species remain to be studied."

**BREEDING SEASON.** Ortmann (1919:174) reported gravid females from February to April.

**HOST FISH.** Unknown (Watters 1994).

**STATUS.** Federally endangered





Map 3      Distribution of *Alasmidonta marginata susquehannae* Ortmann, 1919 in Maryland. Not reported historically from Maryland.

***Alasmidonta marginata susquehannae* Ortmann, 1919** *Susquehanna elktoe* Fig. 5**SYNONYMY.**

*Alasmidonta marginata* Say, 1818  
*Alasmidonta marginata* Say, 1818  
*Mya rugulosa* Wood, 1828  
*Alasmodon (Decurambis) scriptum* Rafinesque, 1831  
*Unio swanaonensis* Hanley, 1842  
*Alasmidonta corrugata* DeKay, 1843  
*Margaritana marginata* "var. *truncata*" B.H. Wright, 1898  
*Alasmidonta (Decurambis) marginata susquehannae*  
 Ortmann, 1919  
*Alasmidonta marginata variabilis* F.C. Baker, 1928  
*Alasmidonta (Decurambis) marginata* Say, 1818



**SHELL DESCRIPTION.** Shell elongated, dorsal margin gently rounded, ventral margin nearly straight, anterior end rounded, shell inflated, relatively thin, posterior ridge sharp and prominent, posterior slope broad, flat and often covered with fine flutings beak sculpture heavy double-looped ridges, periostracum yellowish brown often a rusty color, numerous green rays with dark green spots, pseudocardinal teeth elongate, lateral teeth reduced to a swelling, beak cavity open and moderately deep, nacre color bluish white often with a tinge of salmon.

**DISTRIBUTION:** Maryland, New York, Pennsylvania, (Williams et al. 1993)


**ECOLOGY.** Ortmann (1919:186) reported the ecology of *Alasmidonta marginata* from the Interior Basin as "most decidedly a species of the riffles, being found there in finer or coarse, but firmly packed gravel in swift currents. ... Call (1900) remarks on the strongly developed foot of this species ... When fully extended, the foot is firmly attached to the gravel in the river bed, and it requires quite an effort to dislodge the specimens." He noted that the subspecies in the Susquehanna River Basin "frequents riffles, and this renders it the more conspicuous, since there are few species on the Atlantic side which prefer this habitat." (Ortmann 1919:190).

**BREEDING SEASON.** Ortmann (1919:188) reported eggs and glochidia in August and noted that the species is probably bradyctictic.

**HOST FISH.** Watters (1994:97) lists the following species as host fish for *Alasmidonta marginata*: northern hogsucker, *Hypentelium nigricans*; shorthead redhorese, *Moxostoma macrolepidotum*; white sucker, *Catostomus commersoni*, rock bass, *Ambloplites rupestris*, warmouth, *Lepomis gulosus*. No mention is made of the host fish for the Atlantic Slope subspecies.

**STATUS.** Special concern (Williams et al. 1993).

**COMMENTS:** Gerberich (1984) noted that this taxon has not been reported from Maryland, but it probably occurred historically in the main channel of the Susquehanna River in Maryland.



Map 4

Distribution of *Alasmidonta undulata* (Say, 1817) in Maryland.

***Alasmidonta undulata* (Say, 1817) triangle floater Fig. 6.****SYNONYMY**


*Unio undulata* Say, 1817  
"*Unio glabratus*? Lamarck" Sowerby, 1823  
*Unio hians* Valenciennes, 1827  
*Alasmidonta sculptilis* Say, 1829  
*Uniopsis radiata* Swainson, 1840  
*Uniopsis mytiloides* Swainson, 1840  
*Margaritana triangulata* Lea, 1858  
*Unio swainsoni* Sowerby, 1868  
*Alasmidonta (Alasmidonta) undulata* (Say, 1817)

**SHELL DESCRIPTION.** Shell ovate to triangular, shell thickened and strong anteriorly becoming thin posteriorly, ventral margin evenly rounded, posterior margin rounded and pointed ventrally, beak somewhat inflated and located in the anterior third of the shell, posterior ridge well marked and rounded, posterior slope of adults with a few low coarse ridges, periostracum smooth and shiny, yellowish with green rays, background becoming brownish to black in adult specimens, pseudocardinal teeth stumpy, interdental projection well developed in left valve, nacre white and some with salmon or pink.

**DISTRIBUTION:** Alabama, Connecticut, Delaware, Florida, Georgia, Massachusetts, Maryland, Maine, North Carolina, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, South Carolina, Virginia, Vermont, West Virginia, New Brunswick, Nova Scotia, Ontario, Quebec, Canada (Williams et al. 1993).

**ECOLOGY.** Ortmann (1919:180) noted this species "it is quite evident that it avoids the larger rivers, and prefers the smaller streams, where it becomes locally very abundant, going far up towards the headwaters, ... It does not seem to favor riffles and very rough water, but is found chiefly in more quiet parts, but with some current, for instance, above riffles, where a steady flow of water prevails. It does not like slackwater, but occasional it is found in ponds and canals; .. It also like mill-races, if the current is not too rapid. It lives mostly in a mixture of coarser or finer gravel with sand and mud; but I have taken it also in eddies with slow current embedded in the mud deposited between larger stones."


**BREEDING SEASON.** Ortmann (1919:178) reported gravid females July to September and April to June, a bradyctictic species.



**HOST FISH.** Unknown (Watters 1994).

**STATUS.** Special concern (Williams et al. 1993).





Map 5      Distribution of *Alasmidonta varicosa* (Lamarck, 1819) in Maryland.



***Alasmidonta varicosa* (Lamarck, 1819) brook floater Fig. 7.****SYNONYMY.**

*Unio varicosa* Lamarck, 1819

*Alasmodon corrugata* DeKay, 1843

*Alasmidonta (Decurambis) varicosa* (Lamarck, 1819)

*Mya rugulosa* Wood, 1856

**SHELL DESCRIPTION.** Shell kidney shaped, thin shelled, slightly thicker anteriorly, ventral margin straight and slightly concave centrally, beaks narrow and bluntly pointed, posterior ridge broad, flat and inflated, posterior slope flattened to slightly concave, often with poorly developed corrugations, periostracum yellowish and partly to completely covered with green rays, hinge teeth poorly developed to rudimentary, interdental projection well defined or a small swelling, beak cavity open and shallow, nacre bluish to bluish white, tan to olive or pinkish in the beak cavity.

**DISTRIBUTION:** Connecticut, Delaware, Georgia, Massachusetts, Maryland, Maine, North Carolina, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, South Carolina, Virginia, Vermont, West Virginia, New Brunswick, Nova Scotia, Canada (Williams et al. 1993).


**ECOLOGY.** Ortmann (1919:193) noted " In the smaller streams this species is more abundant, and locally common. ... it is distinctly a shell which prefers strong currents and gravelly bottoms, thus being most frequently found in and near riffles ... it goes far up into the headwaters."

**BREEDING SEASON.** Ortmann (1919:191-192) observed eggs in the marsupium in August, glochidia in September and discharge of glochidia in May. The species is bradytictic.

**HOST FISH.** Unknown (Watters 1994).

**STATUS.** Threatened (Williams et al. 1993).





Map 6

Distribution of *Anodonta implicata* Say, 1829 in Maryland.

***Anodonta implicata* Say, 1829 alewife floater Fig. 8.****SYNONYMY.**

*Anodonta implicata* Say, 1829

*Anodonta newtonensis* Lea, 1836

*Anodonta housatonica* Linsley, 1845

*Anodonta (Pyganodon) implicata* Say, 1829

**SHELL DESCRIPTION.** Shell elongate-elliptical to elongate ovate, valves quite inflated, subcylindrical, valves thick for *Anodonta*, ventral margin straight, dorsal margin straight, posterior ridge rounded and biangulate, umbos somewhat swollen and located in anterior third of shell, beak sculpture is 5 to 7 double looped bars, periostracum smooth, yellowish brown, greenish brown to reddish brown and black with age, immature specimens have fine rays, no teeth, beak cavity shallow, shell distinctly thickened along the anterior ventral margin below the pallial line, nacre color pale copper, pinkish or white.


**STATUS.** Currently stable (Williams et al. 1993).


**DISTRIBUTION:** Connecticut, Delaware, Massachusetts, Maryland, Maine, North Carolina, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, South Carolina, Virginia, Vermont, New Brunswick, Nova Scotia, Quebec, Canada (Williams et al. 1993).

**ECOLOGY.** Ortmann (1919:162) reported the ecology of this species as: "Generally it is reported as living in ponds, but sometimes also in rivers."

**BREEDING SEASON.** Ortmann (1919:160) reported the breeding season as September to May, a bradytic species.

**HOST FISH.** Watters (1994:101) lists the following species as host fish for *Anodonta implicata*: alewife, *Alosa pseudoharengus* (Wilson, 1811).





Map 7      Distribution of *Elliptio angustata* (Lea, 1831) in Maryland.

***Elliptio angustata* (Lea, 1831) Carolina lance fig. 9****SYNONYMY.**

*Unio angustatus* Lea, 1831

**SHELL DESCRIPTION.** The shell is elongated, subrhomboid or subelliptical, subcompressed to slightly inflated, the valves are thin to subsolid, the beaks are scarcely inflated or elevated and their sculpture is a series of strong ridges nearly parallel with the growth lines. The posterior ridge is well developed, angulated and more or less double. Posterior margin bluntly pointed and ventral margin straight. The epidermis is olive in juveniles becoming nearly black in older individuals and the surface is rather dull. The pseudocardinal teeth are compressed and double in both valves, the upper one in the right valve is small. The lateral teeth are long, somewhat club-shaped. Adductor muscle scars are shallow. Nacre color is purplish brown (after Simpson 1914:678).

**DISTRIBUTION:** Georgia, Maryland, North Carolina, South Carolina, Virginia (Williams et al. 1993).

**ECOLOGY.** Ortmann (1919:113) noted "I found it either in gravel, or among larger rocks, in mud-filled interstices, but generally not in strong current, but in quiet coves and eddies. It also seems to prefer the smaller streams to the larger rivers."


**BREEDING SEASON.** Ortmann (1919:111) listed gravid females from early May to early June (this species listed as *Elliptio cupreus*).

**HOST FISH.** Unknown (Watters 1994).

**STATUS.** Special concern (Williams et al. 1993).

**COMMENTS.** There is much confusion regarding the lanceolate *Elliptio* of the middle Atlantic Slope drainages. Lea described numerous species which were subsequently all lumped under *Elliptio lanceolata* (Lea, 1828). Two names have been used for the second lanceolate species *Elliptio producta* (Lea, 1836) described from the Savannah River in Georgia and *Elliptio angustata* (Lea, 1831) described from the Cooper River in South Carolina.





Map 8

Distribution of *Elliptio complanata* (Lightfoot, 1786) in Maryland.

***Elliptio complanata* (Lightfoot, 1786) eastern elliptio Fig. 10****SYNONYMY.**

*Mya complanata* Lightfoot, 1786  
*Unio violaceus* Spengler, 1793  
*Unio purpureus* Say, 1817  
*Unio rarisulcata* Lamarck, 1819  
*Unio coarctata* Lamarck, 1819  
*Unio purpurascens* Lamarck, 1819  
*Unio rhombula* Lamarck, 1819  
*Unio carinifera* Lamarck, 1819  
*Unio georgina* Lamarck, 1819  
*Unio glabrata* Lamarck, 1819  
*Unio sulcidens* Lamarck, 1819  
*Unio virginiana* Lamarck, 1819  
*Unio aurata* Rafinesque, 1820  
*Unio fluviatilis* Green, 1827  
*Mya rigida* Wood, 1828  
*Unio griffithianus* Lea, 1834  
*Unio complanatus subinflatus* Conrad, 1835  
*Unio jejunos* Lea, 1838  
*Unio fuliginosus* Lea, 1845  
*Unio cuvierianus* Lea, 1852  
*Unio errans* Lea, 1856 good species ???  
*Unio vicinus* Lea, 1856  
*Unio geminus* Lea, 1856  
*Unio abbevilensis* Lea, 1857  
*Unio percoarctatus* Lea, 1857  
*Unio wheatleyi* Lea, 1857  
*Unio catawbensis* Lea, 1861  
*Unio insulsus* Lea, 1857  
*Unio spadiceus* Lea, 1857  
*Unio macer* Lea, 1857  
*Unio contractus* Lea, 1857  
*Unio virens* Lea, 1857  
*Unio savannahensis* Lea, 1857  
*Unio subflavus* Lea, 1857  
*Unio fumatus* Lea, 1857  
*Unio subniger* Lea, 1857  
*Unio neusensis* Lea, 1857  
*Unio purus* Lea, 1858  
*Unio exactus* Lea, 1858  
*Unio postellii* Lea, 1858  
*Unio roswellensis* Lea, 1859  
*Unio burkensis* Lea, 1859  
*Unio hallenbeckii* Lea, 1859  
*Unio baldwinensis* Lea, 1859  
*Unio salebrosus* Lea, 1859  
*Unio raeensis* Lea, 1859  
*Unio latus* Lea, 1859  
*Unio quadratus* Lea, 1859  
*Unio squameus* Lea, 1861  
*Unio rostrum* Lea, 1861  
*Unio northamptonensis* Lea, 1861  
*Unio decumbens* Lea, 1861  
*Unio raleighensis* Lea, 1863  
*Unio aberrans* Lea, 1863  
*Unio weldonensis* Lea, 1863  
*Unio mecklenbergensis* Lea, 1863  
*Unio chathamensis* Lea, 1863  
*Unio gastonensis* Lea, 1863  
*Unio quadrilaterus* Lea, 1863  
*Unio indefinitus* Lea, 1863  
*Unio indefinitus* Lea, 1866  
*Unio mediocris* Lea, 1863  
*Unio perlucens* Lea, 1863  
*Unio curatus* Lea, 1863  
*Unio protensus* Lea, 1865  
*Unio lazarus* Sowerby, 1868 in Reeve.  
*Unio beaverensis* Lea, 1868  
*Unio nubilus* Lea, 1868  
*Unio datus* Lea, 1868  
*Unio humerosus* Lea, 1868  
*Unio uhareensis* Lea, 1868  
*Unio tortuosus* Sowerby, 1868

*Unio santeensis* Lea, 1871  
*Unio yadkinensis* Lea, 1872  
*Unio amplius* Lea, 1872  
*Unio ligatus* Lea, 1872  
*Unio differtus* Lea, 1872  
*Unio subparallelus* Lea, 1872  
*Unio oblongus* Lea, 1872  
*Unio curvatus* Lea, 1872  
*Unio irwinensis* Lea, 1872  
*Unio subsquamosus* Lea, 1872  
*Unio infuscus* Lea, 1872  
*Unio ratus* Lea, 1872  
*Unio basalis* Lea, 1872  
*Unio dissimilis* Lea, 1872  
*Unio cirratus* Lea, 1874  
*Unio subolivaceus* Lea, 1874  
*Unio infulgens* Lea, 1874  
*Unio corneus* Lea, 1874  
*Unio dooleyensis* Lea, 1874  
*Unio gesnerii* Lea, 1874  
*Unio invenustus* Lea, 1874  
*Unio (Arconaia) provancheriana* Pilsbry, 1890  
*Unio palliatus* 'Ravenel' Simpson, 1900  
*Unio pullatus majusculus* De Gregorio, 1914  
*Unio complanatus mainensis* Rich, 1915  
*Elliptio (Elliptio) complanata* (Lightfoot, 1786)

**SHELL DESCRIPTION.** Shell outline long, trapezoidal, to rhomboid or subelliptical, shell compressed to inflated and thin to solid, dorsal and ventral margins roughly parallel, ventral margin often straight, posterior ridge broad, double and rounded to angular, beaks low and uninflated, periostracum brownish or yellowish green becoming almost black with age often with green rays over the entire shell, lateral teeth straight, beak cavity very shallow, nacre usually purple, but white to light orange or salmon occur.





**DISTRIBUTION:** Alabama, Connecticut, Delaware, Florida, Georgia, Massachusetts, Maryland, Maine, Michigan, Minnesota, North Carolina, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, South Carolina, Virginia, Vermont, Wisconsin, West Virginia, New Brunswick, Nova Scotia, Ontario, Quebec, Canada (Williams et al. 1993).

**ECOLOGY.** Ortmann (1919:109) recorded the ecology of *Elliptio complanata* as: "It apparently has no ecological preferences, being found practically in any permanent body of water; in canals and reservoirs with quiet water and muddy bottom, as well as in large rivers with strong current and heavy gravel and rocks. In the small creeks it goes very far into the headwaters."


**BREEDING SEASON.** Ortmann (1919:104) reported gravid females from April to mid-July with glochidia present in early June and as late as August. This is a tachytictic species.

**HOST FISH.** Watters (1994:103) lists the following species as host fish for *Elliptio complanata*: banded killifish, *Fundulus diaphanus*; green sunfish, *Lepomis cyanellus*; orangespotted sunfish, *Lepomis humilis*; largemouth bass, *Micropterus salmoides*; white crappie, *Pomoxis annularis*; yellow perch, *Perca flavescens*.

**STATUS.** Currently stable (Williams et al. 1993).



**Page left blank on purpose.**



Map 9      Distribution map of *Elliptio fisheriana* (Lea, 1838) in Maryland.

***Elliptio fisheriana* (Lea, 1838) northern lance Fig. 11.****SYNONYMY.**

*Unio fisherianus* Lea, 1838

*Margarita (Unio) fisherianus* (Lea, 1838)

*Margaron (Unio) fisherianus* (Lea, 1838)

*Elliptio fisherianus* (Lea, 1838)

**SHELL DESCRIPTION.** Shell elongate, compressed, shell thin, beaks low, posterior ridge subangular, posterior end bluntly rounded, ventral margin broadly curved, anterior end rounded, periostracum olive, green to brown and black, often with faint rays, beak cavity very shallow, nacre white to purplish.

**DISTRIBUTION:** Delaware, Maryland, Pennsylvania, Virginia (Williams et al. 1993).


**ECOLOGY.** Johnson (1970:335) lists the ecology of *Elliptio lanceolata* sensu lato as "Lives on sandy bottoms, often found crawling about with much of the shell out of the sand, also found among rocks and in mud, where the current is not too swift."


**BREEDING SEASON.** Ortmann (1919:113) listed the breeding season of this species as unknown.

**HOST FISH.** Unknown (Watters 1994).

**STATUS.** Special concern (Williams et al. 1993).

**COMMENTS.** Type locality for this species is the "head of the Chester River [Kent Co.] Maryland (Johnson 1970:333).






Map 10      Distribution of *Lampsilis cardium* Rafinesque, 1820 in Maryland.

***Lampsilis cardium* Rafinesque, 1820 plain pocketbook Fig. 12.****SYNONYMY.**

*Lampsilis cardium* Rafinesque, 1820  
*Unio ventricosus* Barnes, 1823  
*Unio occidens* Call, 1887  
*Lampsilis ventricosus* (Barnes, 1823)  
*Lampsilis ovata ventricosa* (Barnes, 1823)  
*Unio occidens* Lea, 1829  
*Unio subovatus* Lea, 1831  
*Unio lenis* Conrad, 1838  
*Unio canadensis* Lea, 1857  
*Unio latissimus* Sowerby, 1868  
*Lampsilis ventricosa* var. *lurida* Simpson, 1914  
*Lampsilis ventricosa cohongoronta* Ortmann, 1912  
*Lampsilis ventricosa winnebagoensis* Baker, 1928  
*Lampsilis ventricosa pergloboas* Baker, 1928  
*Lampsilis ovata ventricosa* (Barnes, 1823)



**SHELL DESCRIPTION.** Shell globose, elliptical, female shells often more rectangular, ventral margin evenly rounded, shell relatively thin but stout, moderately inflated, posterior ridge not prominent and rounded, periostracum yellowish to tan becoming darker with age, broad to narrow rays usually cover the shell, some shells may not have rays, pseudocardinal teeth compressed, lateral teeth short, interdentum narrow, beak cavity wide and deep, nacre white. compare with *Lampsilis ovata*.

**DISTRIBUTION:** Alabama, Arkansas, Iowa, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maryland, Michigan, Minnesota, Missouri, Mississippi, Ohio, Oklahoma, Tennessee, Texas, West Virginia, Wisconsin (Williams et al. 1993).


**ECOLOGY.** Ortmann (1919:306) listed the ecology of the species as favoring "rather quiet pools and eddies above and below riffles. Open pools in riffles, in *Dianthera*-patches, with moderate current, and a bottom of fine gravel covered with a thin layer of mud ... occasionally this form is found even in pure sand and rather deep and soft mud."

**BREEDING SEASON.** Ortmann (1919:303) reports gravid females from the end of July to October and May to early July. This species is bradyctictic.

**HOST FISH.** Watters (1994:108) lists the following species as host fish for *Lampsilis cardium*: bluegill, *Lepomis macrochirus*; largemouth bass, *Micropterus salmoides*; smallmouth bass, *Micropterus dolomieu*; white crappie, *Pomoxis annularis*; sauger, *Stizostedion canadense*, walleye, *Stizostedion vitreum*; yellow perch, *Perca flavescens*.

**STATUS.** Special concern (Williams et al. 1993).

**COMMENTS.** The figure specimen is the lectotype of *Lampsilis ventricosa cohongoronta* Ortmann, 1912, CMNH 61.3994. The type locality of this subspecies is the Potomac River at Hancock, Washington County, Maryland.



Map 11      Distribution of *Lampsilis cariosa* (Say, 1817) in Maryland

***Lampsilis cariosa* (Say, 1817) yellow lampmussel Fig. 13.****SYNONYMY.**

*Unio cariosus* Say, 1817

*Margarita (Unio) cariosus* (Say, 1817)

*Margaron (Unio) cariosus* (Say, 1817)

*Lampsilis pallida* Rafinesque, 1820

*Unio ovata* Valenciennes, 1827 PREOCCUPIED

*Unio viridis* Ferussac, 1835

*Unio crocatus* Lea, 1841

*Unio oratus* Conrad, 1849

*Lampsilis cariosa* (Say, 1817)

*Lampsilis (Lampsilis) cariosa* (Say, 1817)



**SHELL DESCRIPTION.** Shell medium size, male shell elliptical and somewhat elongate, female shells subobovate to obovate, moderately inflated, thick, anterior margin rounded, ventral margin slightly curved, dorsal margin straight, posterior ridge rounded, umbos swollen, and raised just above the hinge line and located anterior of the midline of the shell, periostracum shiny, waxy or straw yellow, becoming darker brownish yellow, green or black rays when present usually restricted to the posterior slope, pseudocardinal teeth compressed, beak cavity moderately deep, nacre color white or tinged with salmon.

**DISTRIBUTION:** Connecticut, Delaware, Georgia, Massachusetts, Maine, Maryland, North Carolina, New Jersey, New York, Pennsylvania, South Carolina, Virginia, New Brunswick, Nova Scotia, Canada (Williams et al. 1993).


**ECOLOGY.** Ortmann (1919:317) reported "Where found this species is generally abundant, even in smaller streams ... It is always found in lively currents, on shoals and riffles, in finer or coarser gravel, and very often in bars of pure sand."

**BREEDING SEASON.** Ortmann (1919:315) was only able to establish gravid females in August, with glochidia appearing in late August. This is a bradyctictic species releasing glochidia in the spring or early summer.

**HOST FISH.** Unknown (Watters 1994).

**STATUS.** Threatened (Williams et al. 1993). Maryland DNR listed this species as extinct in Maryland (Anonymous, 1994).






Map 12      Distribution of *Lampsilis radiata* (Gmelin, 1791) in Maryland

***Lampsilis radiata* (Gmelin, 1791) eastern lampmussel Fig. 14.****SYNONYMY.**

*Mya radiata* Gmelin, 1791  
*Unio luteola* Lamarck, 1819  
*Unio lineata* 'Valenciennes' Bory de St. Vincent, 1827  
*Unio tenebrosus* Conrad, 1834  
*Unio melinus* Conrad, 1838  
*Unio boydianus* Lea, 1840  
*Unio rosaceus* De Kay, 1843  
*Mya oblongata* Wood, 1856  
*Unio elongata* S.G. Goodrich, 1858  
*Unio obliquiradiatus* Reeve, 1865  
*Unio conspicuus* Lea, 1872  
*Unio virginiana* Simpson, 1900  
*Lampsilis radiata* (Gmelin, 1791)  
*Lampsilis radiata oneidensis* Baker, 1916  
*Unio virginea* Frierson, 1927  
*Lampsilis radiata radiata* (Gmelin, 1791)  
*Lampsilis radiata* var. *conspicua* (Lea, 1872)  
*Lampsilis* (*Lampsilis*) *radiata radiata* (Gmelin, 1791)



**SHELL DESCRIPTION.** Shell medium to large, subelliptical or subovate in outline, moderately elongate, valves from not inflated to quite inflated, and solid, anterior end rounded, posterior end in females broadly expanded and rounded, dorsal margin straight and ventral margin straight to gently curved, posterior ridge mostly absent, posterior slope broad, beaks rather sharp but not very inflated, periostracum yellowish or brownish green with dark green rays over the entire surface, no interdentum, beak cavity shallow, nacre white, may be tinged with pink or salmon.


**DISTRIBUTION:** Connecticut, Delaware, Massachusetts, Maryland, Maine, Michigan, North Carolina, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, South Carolina, Virginia, Vermont, West Virginia, New Brunswick, Nova Scotia, Ontario, Quebec, Canada (Williams et al. 1993).

**ECOLOGY.** Ortmann (1919:317) recorded this species "Where found, this species is generally abundant, even in smaller streams ... It is always found in lively current, on shoals and riffle, in finer or coarser gravel, and very often in bars of pure sand."

**BREEDING SEASON.** Ortmann (1919:293) only reported gravid females in late August and glochidia in gravid females in late August.

**HOST FISH.** Watters (1994:107) lists the following species as host fish for *Lampsilis radiata*: rockbass, *Ambloplites rupestris*; pumpkinseed, *Lepomis gibbosus*; largemouth bass, *Micropterus salmoides*; smallmouth bass, *Micropterus dolomieu*; black crappie, *Pomoxis nigromaculatus*; yellow perch, *Perca flavescens*.

**STATUS.** Currently stable (Williams et al. 1993).



Map 13      Distribution of *Lasmigona subviridis* (Conrad, 1835) in Maryland.

***Lasmigona subviridis* (Conrad, 1835) green floater Fig. 15.****SYNONYMY.**

*Unio subviridis* Conrad, 1835


*Unio tappanianus* Lea, 1838

*Unio hyalinus* Lea, 1845

*Margaritana quadrata* Lea, 1861

*Unio pertenus* Lea, 1863

*Lasmigona (Platynaias) subviridis* (Conrad, 1835)



**SHELL DESCRIPTION.** Shell ovate trapezoid, thin and rather fragile, beaks only projecting slightly above the hinge line, posterior ridge rounded, periostracum light yellow or brown with numerous green rays especially in juveniles, pseudocardinal and lateral teeth small and delicate, beak cavity shallow, nacre whitish to bluish and iridescent posteriorly.


**DISTRIBUTION.** Georgia, Kentucky, Maryland, New Jersey, New York, North Carolina, Pennsylvania, South Carolina, Tennessee, Virginia, West Virginia (Williams et al. 1993).

**ECOLOGY.** Ortmann (1919:124) listed the ecology of this species: "this species is very erratic in its distribution. ... The specimens found by myself in larger rivers generally were few, and often in small branches of the river. But even in small streams, it is not everywhere present. ... it is averse to very strong current, and prefers more quiet parts, pools or eddies with gravelly and sandy bottoms, and it also goes into canals."

**BREEDING SEASON.** Ortmann (1919:122) reported gravid females from August to September and April to June, a bradytictic species.

**HOST FISH.** Unknown (Watters 1994).

**STATUS.** Threatened (Williams et al. 1993).



Map 14      Distribution of *Leptodea ochracea* (Say, 1817) in Maryland.

***Leptodea ochracea* (Say, 1817) tidewater mucket Fig. 16.****SYNONYMY.**

*Mytilus fluviatilis* Gmelin 1791 (Nomen dubium)

*Unio ochraceus* Say, 1817


*Lampsilis rosea* Rafinesque, 1820

*Unio rosaceus* Conrad, 1849

*Lampsilis ochracea* (Say, 1817)

*Lampsilis (Lampsilis) ochracea* (Say, 1817)

*Leptodea ochracea* (Say, 1817)



**SHELL DESCRIPTION.** Shell relatively small, male shell elliptical and female more ovate, shell subinflated and thin, and strong, anterior end rounded, posterior margin evenly rounded, somewhat pointed in males and truncated in females, posterior ridge rounded, often with a few ridges or wrinkles, beaks moderately swollen and raised above the hinge line, located near the middle of the shell, periostracum slightly shiny, brownish olive, greenish-yellow, yellow or reddish yellow, often with fine green rays, over most of the shell, pseudocardinal teeth compressed, lateral teeth short and curved, beak cavities shallow, nacre white or pinkish.


**DISTRIBUTION:** Connecticut, Delaware, Georgia, Massachusetts, Maryland, Maine, North Carolina, New Jersey, New York, Pennsylvania, Rhode Island, South Carolina, Virginia, New Brunswick, Nova Scotia, Canada (Williams et al. 1993).

**ECOLOGY.** Ortmann (1919:320) noted *Leptodea ochracea* was found in "most tide waters north of the Savannah River. ... It is a form of estuaries, ponds, canals, and ditches, probably with more or less muddy bottoms."

**BREEDING SEASON.** Ortmann (1919:319) noted that Isaac Lea had reported this species was gravid in the autumn (October, November). There is a lack of information on the anatomy and biology of this species.

**HOST FISH.** Unknown (Watters 1994).

**STATUS.** Special concern (Williams et al. 1993).



Map 15      Distribution of *Ligumia nasuta* (Say, 1817) in Maryland.



***Ligumia nasuta* (Say, 1817) eastern pondmussel Fig. 17.****SYNONYMY.**

*Unio nasutus* Say, 1817

*Obliquaria attenuata* Rafinesque, 1820

*Unio rostrata* Valenciennes, 1827


*Unio vancouverianus* Sowerby, 1868

*Unio fisherianus* Kuester, 1860 non Lea, 1838

*Lampsilis nasuta* (Say, 1817)

*Eurynia nasuta* (Say, 1817)

*Ligumia nasuta* (Say, 1817)



**SHELL DESCRIPTION.** Shell elongated, subelliptical, shell is over twice as long as high, shell rather thin, subinflated, posterior end bluntly pointed, dorsal margin straight, ventral margin curved, posterior ridge distinct, beaks low and located in anterior quarter of the shell, periostracum greenish yellow, to dark olive or brown sometimes with distinct narrow rays, beak cavity shallow, lateral teeth long and straight, pseudocardinal teeth compressed, nacre white. Male shells taper to a blunt point posteriorly, female shell is distinctly swollen posteriorly.


**DISTRIBUTION:** Connecticut, Delaware, Massachusetts, Maryland, Maine, Michigan, North Carolina, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, South Carolina, Virginia, Ontario, Canada (Williams et al. 1993).

**ECOLOGY.** Ortmann (1919:275) recorded the ecology of this species as "Sandy bottom of great, quiet bodies of water (tidewaters, lakes and probably also canals) seem to furnish the conditions most favorable to this species."

**BREEDING SEASON.** Ortmann (1919: reported gravid females from eastern Pennsylvania from September and May. He noted gravid females from Lake Erie from August to early July, a bradytictic species.

**HOST FISH.** Unknown (Watters 1994).


**STATUS.** Special concern (Williams et al. 1993).



Map 16      Distribution of *Pyganodon cataracta* (Say, 1817) in Maryland.

***Pyganodon* (= *Anodonta*) *cataracta* (Say, 1817) eastern floater Fig. 18.****SYNONYMY.**

*Anodonta cataracta* Say, 1817  
*Anodonta marginata* Say, 1817  
*Anodonta teres* Conrad, 1834  
*Anodon excurvata* De Kay, 1843  
*Anodonta virgulata* Lea, 1857  
*Anodonta lacustris* Lea, 1857  
*Anodonta hallenbeckii* Lea, 1858  
*Anodonta gesnerii* Lea, 1858  
*Anodonta dariensis* Lea, 1858  
*Anodonta williamsii* Lea, 1862  
*Anodonta tryoni* Lea, 1862  
*Anodonta dolearis* Lea, 1863  
*Anodonta doliaris* Lea, 1866  
*Anodonta* (*Pyganodon*) *cataracta cataracta* Say, 1817



**SHELL DESCRIPTION.** Shell medium to large, outline subelliptical and elongate, shells of juveniles not much inflated but much inflated in adult shells, shell thin, posterior margin pointed, ventral margin straight to slightly curved, posterior ridge indistinct, posterior slope often with two faint ridges, beaks slightly swollen, located in anterior third of shell, periostracum usually smooth, shiny, straw yellow to light green grading to a dark green, rays on posterior slope darker, rays on disk of the shell, no teeth are present, muscle scars poorly defined, nacre bluish white.


**DISTRIBUTION:** Alabama, Connecticut, Delaware, Georgia, Massachusetts, Maryland, Maine, Michigan, North Carolina, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, South Carolina, Virginia, Vermont, Wisconsin, West Virginia, New Brunswick, Nova Scotia, Ontario, PE, Quebec, Canada (Williams et al. 1993).

**ECOLOGY.** Ortmann (1919:158) noted a creek and a pond form of this species and remarked about the species: "I collected the pond-form in rather small (artificial) ponds, with very muddy bottoms, and also, but more rarely, in quiet pools in creeks. ... the creek-form ... It seems to avoid the larger rivers ... It favors smaller rivers and creeks. Here it lives on gravelly bottoms, in more or less strongly flowing water (even in riffles), or in more quiet pools in gravel, sand, or mud. The short high form ... seems to prefer large rivers with muddy bottoms."

**BREEDING SEASON.** Ortmann (1919:153) reported gravid females from late July to late April and noted the species is listed as gravid until May, a bradyctictic species.

**HOST FISH.** Watters (1994:99) lists the following species as host fish for *Pyganodon cataracta* as: carp, *Cyprinus carpio*; pumpkinseed, *Lepomis gibbosus*; threespine stickleback, *Gasterosteus aculeatus*; white sucker, *Catostomus commersoni*.

**STATUS.** Currently stable (Williams et al. 1993).




Map 17      Distribution of *Strophitus undulatus* (Say, 1817) in Maryland.

***Strophitus undulatus* (Say, 1817) squawfoot Fig. 19.****SYNONYMY.**

*Anodonta undulata* Say, 1817  
*Alasmodonta edentula* Say, 1820  
*Anodon rugosus* Swainson, 1822  
*Anodonta arkansensis* Call, 1885  
*Anodonta edentula* (Say, 1817)  
*Strophitus edentulus* (Say, 1817)  
*Strophitus rugosus* (Swainson)  
*Anodonta pennsylvanica* [sic] Lamarck, 1819  
*Anodon areolatus* Swainson, 1829  
*Alasmodonta edentula* Say, 1829  
*Anodonta virgata* Conrad, 1836  
*Anodonta pavonia* Lea, 1836  
*Anodonta wardiana* Lea, 1838  
*Anodon unadilla* De Kay, 1843  
*Anodonta tetragona* Lea, 1845  
*Anodonta arkansensis* Lea, 1852  
*Anodonta shaefferiana* Lea, 1852  
*Alasmodon rhombica* Anthony, 1865  
*Anodon papyracea* Anthony, 1865  
*Anodon annulatus* Sowerby, 1867  
*Anodon quadriplicatus* Sowerby, 1867  
*Anodonta salmonia* Clessin, 1873  
*Strophitus undulatus ovatus* Frierson, 1927  
*Strophitus rugosus pepinensis* Baker, 1928  
*Strophitus rugosus winnebagoensis* Baker, 1928  
*Strophitus rugosus lacustris* Baker, 1928  
*Strophitus edentulus* (Say, 1817)

**SHELL DESCRIPTION.** Shell oblong oval, dorsal and ventral margins rounded, shell thin and fragile when young but becoming thicker, length to 11 cm. compressed to inflated, posterior ridge rounded, beaks narrow but only slightly raised above the hinge, periostracum yellow-brown to greenish becoming darker with age, green rays may cover the shells. Pseudocardinal teeth are only swellings or thickening along the hinge, lateral teeth absent, beak cavity shallow, hinge line undulate, nacre white with cream color in beak cavity.

**DISTRIBUTION:** Alabama, Arkansas, Colorado, Connecticut, Delaware, Georgia, Iowa, Illinois, Indiana, Kansas, Kentucky, Louisiana, Massachusetts, Maryland, Maine, Michigan, Minnesota, Missouri, Mississippi, North Carolina, North Dakota, Nebraska, New Hampshire, New Jersey, New York, Ohio, Oklahoma, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Virginia, Vermont, Wisconsin, West Virginia, Manitoba, New Brunswick, Nova Scotia, Ontario, Quebec, Saskatchewan, Canada (Williams et al. 1993).




**ECOLOGY.** Ortmann (1919:204-205) reported "it is distinctly averse to large rivers. ... We may ... call *S. edentulus* a form characteristic of smaller streams ... Baker ... says indeed that it is found in the larger lakes and rivers on muddy bottoms, while Scammon ... reports it to prefer mud and quiet water, and to be the most abundant in small streams (in Kansas). ... in the small streams it avoids riffles, but delights in quiet and protected nooks, pools, and eddies, where there is a moderate and rather uniform current, and a deposit of fine gravel, or sand."

**BREEDING SEASON.** Ortmann (1919:196) reported that the Atlantic slope individuals were "gravid in December and March, but not in April and May." He noted that the Ohio River basin populations (listed as *Strophitus edentulus*) were gravid from July to November and April to May, a typical bradytictic species (Ortmann 1919:198).

**HOST FISH.** Watters (1994:113) lists the following species as host fish for *Strophitus undulatus* as: creek chub, *Semotilus atromaculatus*; largemouth bass, *Micropterus salmoides*. Other authors have noted that no host fish is needed or have listed "several fish".

**STATUS.** Currently stable (Williams et al. 1993).



Map 18      Distribution of *Utterbackia imbecillis* (Say, 1829) in Maryland.

***Utterbackia* (= *Anodonta*) *imbecillis* Say, 1829 paper pondshell Fig. 20.****SYNONYMY.***Anodonta imbecillis* Say, 1829*Anodonta imbecilis* [sic] Say, 1829*Anodonta incert* Lea, 1834*Anodon horda* Gould, 1855*Anodonta henryana* Lea, 1857*Utterbackia imbecillis fusca* Baker, 1927*Anodonta ohioensis* Rafinesque, 1820 [in part]*Anodonta* (*Utterbackia*) *imbecilis* Say, 1829


**SHELL DESCRIPTION.** Shell elongate, oblong, dorsal hinge line rather short and straight, the ventral margin rounded, shell inflated and thin growing to about 10 cm in length, beaks flat, not above the hinge line, beaks on the anterior third of the shell, periostracum shiny, light yellow on the umbo, rest of shell bright green with rays and posterior slope area is dark to black, both valves edentulous, nacre bluish-white, often with pink tinges.

**DISTRIBUTION:** Alabama, Arkansas, Florida, Georgia, Iowa, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maryland, Michigan, Minnesota, Missouri, Mississippi, North Carolina, Nebraska, New Mexico, New York, Ohio, Oklahoma, Pennsylvania, South Carolina, Tennessee, Texas, Virginia, Wisconsin, West Virginia, Ontario, Canada, Mexico (Williams et al. 1993).

**ECOLOGY.** Johnson (1970:364) noted "Lives in soft mud or sand in ponds, creeks, and near the banks of larger rivers."

**BREEDING SEASON.** Ortmann (1919:163) listed this species as bradytictic the females beginning gravid in June and July and discharging glochidia in May noted "the succeeding breeding seasons overlap in June and July, but probably not in the same individual." This species is simultaneous hermaphrodite (Hoeh 1991).

**HOST FISH.** Watters (1994:113) lists the following species as host fish for *Utterbackia imbecillis* as: banded killifish, *Fundulus diaphanus*; creek chub, *Semotilus atromaculatus*; rockbass, *Ambloplites rupestris*; bluegill, *Lepomis macrochirus*; dollar sunfish, *Lepomis marginatus*; green sunfish, *Lepomis cyanellus*; longear sunfish, *Lepomis megalotis*; pumpkinseed, *Lepomis gibbosus*; warmouth, *Lepomis gulosus*; largemouth bass, *Micropterus salmoides*; western mosquitofish, *Gambusia affinis*; yellow perch, *Perca flavescens*.



**STATUS.** Currently stable (Williams et al. 1993).



**Page left blank on purpose.**



Fig. 2 *Dreissena polymorpha* (Pallas, 1771)



Fig. 3 *Corbicula fluminea* (Müller, 1776)



Fig. 4 *Alasmidonta heterodon* (Lea, 1829)



Fig. 5 *Alasmidonta marginata susquehannae* Ortmann, 1919




Fig. 6 *Alasmidonta undulata* (Say, 1817)



Fig. 7 *Alasmidonta varicosa* (Lamarck, 1819)



Fig. 8 *Anodonta implicata* Say, 1829



Fig. 9 *Elliptio angustata* (Lea, 1831)



Fig. 10 *Elliptio complanata* (Lightfoot, 1786)




Fig. 11 *Elliptio fisheriana* (Lea, 1838)



Fig. 12 *Lampsilis cardium* Rafinesque, 1820



Fig. 13 *Lampsilis cariosa* (Say, 1817)





Fig. 14 *Lampsilis radiata* (Gmelin, 1791)



Fig. 16 *Leptodea ochracea* (Say, 1817)



Fig. 18 *Pyganodon cataracta* (Say, 1817)



Fig. 15 *Lasmigona subviridis* (Conrad, 1835)



Fig. 17 *Ligumia nasuta* (Say, 1817)



Fig. 19 *Strophitus undulatus* (Say, 1817)



Fig. 20 *Utterbackia imbecillis* (Say, 1829)

## INTRODUCTION TO THE LITERATURE ON FRESHWATER BIVALVES

## 1. Sphaeriidae

- Burch, J.B. 1975. *Freshwater sphaeriacean clams (Mollusca: Pelecypoda) of North America*. Malacological Publications, Hamburg, Michigan. pp. xi, 1-96

## 2. Corbiculidae

- Clarke, A.H. 1986. Competitive exclusion of *Canthytia* (Unionidae) by *Corbicula fluminea* (Müller). *Malacology Data Net* 1(1):3-10.
- Counts, C.L., III. 1991. *Corbicula* (Bivalvia: Corbiculidae). Part 1. Catalog of fossil and Recent nominal species. Part 2. Compendium of zoogeographic records of North America and Hawaii, 1924-1984. *Tryonia* 21:ii, 1-134
- Gardner, J.A., Jr., W.R. Woodall, Jr., A.A. Staats, Jr. and J.F. Napoli. 1976. The invasion of the Asiatic clam (*Corbicula manilensis* Philippi) in the Altamaha River, Georgia. *The Nautilus* 90(3):117-125.
- Hillis, D.M. and J.C. Patton. 1982. Morphological and electrophoretic evidence for two species of *Corbicula* (Bivalvia: Corbiculidae) in North America. *American Midland Naturalist* 108:74-80.
- Kraemer, L.R. 1979. *Corbicula* (Bivalvia: Sphaeriacea) vs. indigenous mussels (Bivalvia: Unionacea) in U.S. rivers: A hard case for interspecific competition. *American Zoologist* 19(4):1085-1096.
- Leff, L.G., J.L. Burch, and J.V. McArthur. 1990. Spatial distribution, seston removal, and potential competitive interactions of the bivalves *Corbicula fluminea* and *Elliptio complanata* in a coastal plain stream. *Freshwater Biology* 24(2):409-416.
- Potter, J.M. and L.H. Liden. 1986. *Corbicula* control at the Potomac River stream electric station Alexandria, Virginia. *American Malacological Bulletin Special Edition* 2:53-58.
- Rodgers, J.H., Jr., D.S. Cherry, J.R. Clark, K.L. Dickson and J. Cairns, Jr. 1977. The invasion of the Asiatic clam, *Corbicula manilensis* in the New River, Virginia. *The Nautilus* 91(2):43-46.

## 3. Dreissenidae

- Andrusov, N. 1897-1898. Fossile und lebende Dreissensidae Eurasiens. *Travaux de la Société des Naturalistes de St.-Petersbourg. (Section de Géologie et de Minéralogie)* 25: 25 plates (1897); text (1898), 1-687 pp. [In Russian], 1-115 pp. [German resumé].
- Antonov, P.J. 1984. The possibility of age determination in *Dreissena polymorpha* by the structure of the septal surface. *Malacological Review* 17:143 [Abstract].
- Borcherding, J. 1991. The annual reproductive cycle of the freshwater mussel *Dreissena polymorpha* Pallas in lakes. *Oecologia* 87:208-218.
- Eckroat, L.R. and L.M. Steele. 1993. Comparative morphology of the byssi of *Dreissena polymorpha* and *Mytilus edulis*. *American Malacological Bulletin*. 10(1):103-108.
- French, J.R.P. III. 1990. The exotic zebra mussel-a new threat to endangered mussels. *Endangered Species Technical Bulletin* 15(11):3-4.
- Frisina, A.C. and L.R. Eckroat. 1992. Histological and morphological attributes of the byssus of the zebra mussel, *Dreissena polymorpha* (Pallas). *Journal of the Pennsylvania Academy of Science*. 66(2):63-67.
- Garton, D.W. and W.R. Haag. 1991. Heterozygosity, shell length and metabolism in the European mussel, *Dreissena polymorpha*, from a recently established population in Lake Erie. *Comparative Biochemistry and Physiology* 99A:45-48.
- Garton, D.W. and A.M. Stoeckmann. 1991. Genotype-dependent metabolism at the phosphoglucose isomerase locus [sic] at ambient and elevated temperatures. *Proceedings of the Second International Zebra Mussel Research Conference, Rochester New York*, p. 2 [abstract].
- Griffiths, R.W., W.P. Kovalak, and D.W. Schloesser. 1989. The zebra mussel, *Dreissena polymorpha* (Pallas, 1771), in North America: impact on raw water users. IN: *Proceedings: EPRI Service Water System Reliability Improvement Seminar*. Electric Power Research Institute, Palo Alto, CA. pp. 11-27
- Griffiths, R.W., D.W. Schloesser, J.H. Leach, and W.P. Kovalak. 1991. Distribution and dispersal of the zebra mussel (*Dreissena polymorpha*). *Canadian Journal of Fisheries and Aquatic Sciences* 48:1381-1388.
- Haas, F. 1938. Bivalvia. Pp. 209-466 IN: H. G. Bronn's *Klassen und Ordnungen des Tierreichs* 2(2). Akademische Verlagsgesellschaft, Leipzig.
- Hebert, P.D.N., B.W. Muncaster, and G.L. Mackie. 1989. Ecological and genetic studies on *Dreissena polymorpha* (Pallas): a new mollusc in the Great Lakes. *Canadian Journal of Fisheries and Aquatic Sciences* 46:1587-1591.
- Hebert, P.D.N., C.C. Wilson, M.H. Murdoch, and R. Lazar. 1991. Demography and ecological impact of the invading mollusc *Dreissena polymorpha*. *Canadian Journal of Zoology* 69:405-409.
- Hopkins, G.J. 1990. The zebra mussel, *Dreissena polymorpha*: a photographic guide to the identification of microscopic veligers. Available from Queen's Printer for Ontario, ISBN:0-7729-6349-5.
- Hunter, R.D. and J.F. Bailey. 1992. *Dreissena polymorpha* (zebra mussel): colonization of soft substrate and some effects on unionid bivalves. *The Nautilus* 106:60-67.



- Keen, A.M. 1969. Superfamily Dreissenacea. Pp. N643-N644 IN: R. C. Moore (ed.) *Treatise on Invertebrate Paleontology, part N, vol. 2, Mollusca 6, Bivalvia*. Geological Society of America.
- Kennard, A.S. and B.B. Woodward. 1926. *SYNONYMY of the British non-marine Mollusca (Recent and post-Tertiary)*. British Museum, London. xxiv, 447 pp.
- Kerney, M.P. & B.S. Morton. 1970. The distribution of *Dreissena polymorpha* (Pallas) in Britain. *Journal of Conchology* 27:97-100.
- Kornobis, S. 1977. Ecology of *Dreissena polymorpha* (Pall.) (Dreissenidae, Bivalvia) in lakes receiving heated water discharges. *Pol. Arch. Hydrobiol.* 24:531-545
- Leentvaar, P. 1975. Geographical distribution and biology of *Dreissena polymorpha* Pallas. *Hydrobiological Bulletin* 9:120-122.
- Lewandowski, K. 1983. Formation of annuli on shells of young *Dreissena polymorpha* (Pall.). *Pol. Arch. Hydrobiol.* 30:343-351
- Lewandowski, K. 1976. Unionidae as a substratum for *Dreissena polymorpha* Pall. *Pol. Arch. Hydrobiol.* 23:409-420
- Lewandowski, K. 1982. The role of early development stages in the dynamics of *Dreissena polymorpha* (Pall.) (Bivalvia) populations in lakes. II. Settling of larvae and the dynamics of numbers of settled individuals. *Ekol. Pol.* 30:223-286
- Limanova, N.A. 1964. *Dreissena* (Bibliography). pp. 71-145. IN: Shtegman, B.K. (Ed.) *Biology and Control of Dreissena: a collection of papers. Academy of Sciences of the USSR. Institute of Biology of Inland Waters.* 7(10). [In Russian]. [Israel Program for Scientific Translations, 1968].
- Locard, A. 1893. Les *Dreissensia* du système européen d'après la collection Bourguignat. *Revue Suisse de Zoologie* 1:113-185, pls. 5-7.
- Ludyanskiy, M. 1991. Are *Dreissena polymorpha* and *Dreissena bugensis* synonymous? *Dreissena polymorpha Information Review* 2(6):1-3
- L'vova, A.A. and Y.I. Starobogatov. 1982. A new species of the genus *Dreissena* (Bivalvia, Dreissenidae) from the Okhridsky Lake. *Zoologicheskii Zhurnal* 61:1749-1751 [in Russian].
- MacIsaac, H.J. and W.G. Sprules. 1991. Ingestion of small-bodied zooplankton by zebra mussels (*Dreissena polymorpha*): can cannibalism on larvae influence population dynamics? *Canadian Journal of Fisheries and Aquatic Sciences* 48(11):2051-2060.
- Mackie, G.L. 1991. Biology of the exotic zebra mussel, *Dreissena polymorpha*, in relation to native bivalves and its potential impact in Lake St. Clair. *Hydrobiologia* 219:251-268.
- Mackie, G.L., W.N. Gibbons, B.W. Muncaster, and I.M. Gray. 1989. The zebra mussel, *Dreissena polymorpha*: a synthesis of European experiences and a preview for North America, Report prepared for Wat. Resour. Br., Great Lakes Sect. Available from Queen's Printer for Ontario, ISBN 0-7729-5647-2
- McMahan, R.F. and J.L. Tsou. 1990. Impact of European zebra mussel infestation to the electric power industry. *Proceedings of the American Power Conference* 52:988-997
- Marelli, D.[C.] 1991. Zhadin's systematic account of *Dreissena*: are the species valid? *Dreissena polymorpha Information Review* 2(4):3.
- Marelli, D. 1994. The phylogeny of the Dreissenidae: a Phylogenetic reconstruction of the zebra mussel tree. *Dreissenal, Zebra Mussel Information Clearinghouse* 5(4):1
- Marsden, J.E. 1991. Genetics of zebra mussels: Critical data for ecological studies and development of effective long-term control strategies. *Dreissena polymorpha Information Review* 2(5):3-4.
- Marsden, J.E. 1991. Standard protocols for monitoring and sampling zebra mussels. [Draft]. Center for aquatic Ecology, Illinois Natural History Survey. *Aquatic Ecology Technical Report* 91/4:i-v, 1-34.
- Marsden, J.E. and B. May. 1991. Genetic polymorphism among disjunct populations of zebra mussels in the Great Lakes. *Proceedings of the Second International Zebra Mussel Research Conference, Rochester New York*, p. 11 [abstract].
- May, B. and J.E. Marsden. 1992. Genetic identification and implications of another invasive species of dreissenid mussel in the Great Lakes. *Canadian Journal of Fisheries and Aquatic Sciences* 49(7):1501-1506.
- Morton, B. 1969. Studies on the biology of *Dreissena polymorpha* Pall. I. General anatomy and morphology. *Proceedings of the Malacological Society of London* 38:301-321.
- Morton, B. 1969. Studies on the biology of *Dreissena polymorpha* Pall. II. Correlation of the rhythms of adductor activity, feeding, digestion and excretion. *Proceedings of the Malacological Society of London* 38:401-414.
- Morton, B. 1969. Studies on the biology of *Dreissena polymorpha* Pall. III. Population dynamics, *Proceedings of the Malacological Society of London* 38:471-482
- Morton, B. 1969. Studies on the biology of *Dreissena polymorpha* Pall. 4. Habitats, distribution and control. *Water treatment and Examination* 18:233-240.
- Morton, B. 1970. The evolution of the heteromyarian condition in the Dreissenacea (Bivalvia). *Palaeontology* 13:563-572.
- Morton, B. 1971. Studies on the biology of *Dreissena polymorpha* Pall. V. Some aspects of filter feeding and the effect of micro-organisms upon the rate of filtration, *Proceedings of the Malacological Society of London* 39:289-301
- Neary, B.P. and J.H. Leach. 1992. Mapping the potential spread of the zebra mussel (*Dreissena polymorpha*) in Ontario. *Canadian Journal of Fisheries and Aquatic Sciences* 49(2):406-415.

- Nalepa, T.F. and D.W. Schloesser. 1992. *Zebra mussels: Biology, impacts, and control*. Lewis Publishers, Boca Raton. 810 pp.
- Nuttall, C.P. 1990. Review of the Caenozoic heterodont bivalve superfamily Dreissenacea. *Palaeontology* 33:707-737, 6 pls.
- Oldham, C. 1930. Locomotive habit of *Dreissena polymorpha*. *Journal Conchology of London* 19:25-26
- O'Neil, C.R., jr. and D.B. MacNeil. 1989. *Dreissena polymorpha*: an unwelcome new Great Lakes invader. *Sea Grant Fact Sheet*. pp. 1-8.
- Piesik, Z. 1983. Biology of *Dreissena polymorpha* settling of stylon nets and the role of this mollusk in eliminating the seston and the nutrients from the water course. *Pol. Arch. Hydrobiol.* 30:353-362
- Ramchran, C.W., D.K. Padilla, and S.I. Dodson. 1992. A multivariate model for predicting population fluctuations of *Dreissena polymorpha* in North American lakes. *Canadian Journal of Fisheries and Aquatic Sciences* 49(2):150-158.
- Ramchran, C.W., D.K. Padilla, and S.I. Dodson. 1992. Models to predict potential occurrence and density of the zebra mussel, *Dreissena polymorpha*. *Canadian Journal of Fisheries and Aquatic Science*. 49(12):2611-2620.
- Reeders, H.H., A. Bij de Vaate, and F.J. Slim. 1989. The filtration rate of *Dreissena polymorpha* (Bivalvia) in three Dutch lakes with reference to biological water quality management. *Freshwater Biology* 22:133-141
- Rose, J.L. and L. Eckroat. 1991. Genetic composition and characterization of five zebra mussel populations in the Great Lakes. *Proceedings of the Second International Zebra Mussel Research Conference, Rochester New York*, p. 12 [abstract].
- Rosenberg, G. and M.L. Ludyanskiy. 1994. A nomenclatural review of *Dreissena* (Bivalvia: Dreissenidae), with identification of the quagga mussel as *Dreissena bugensis*. *Canadian Journal of Fisheries and Aquatic Sciences* 51(7):1474-1484.
- Scheil, H.-G., and A. Günther. 1981. Amylases in *Dreissena polymorpha* Pall. (Mollusca: Eulamellibranchiata): evidence for two polymorphic systems. *Zoologischer Anzeiger* 207:120-122.
- Schloesser, D.W. and W.P. Kovalak. 1991. Infestation of unionids by *Dreissena polymorpha* in a power plant canal in Lake Erie. *Journal of Shellfish Research*. 10(2):355-359.
- Secchi, F. and M. Martoja. 1986. Existence de populations uricotéliques chez *Abra alba* et *Dreissena polymorpha* (Bivalves Eulamellibranchies). *Canadian Journal of Fisheries and Aquatic Sciences* 43:2504-2508.
- Sigurdson, J.B., C.W. Titman, and P.A. Davies. 1976. The dispersal of young post-larval bivalve molluscs by byssus threads. *Nature* 262:386-387
- Sprung, M. 1987. Ecological requirements of developing *Dreissena polymorpha* eggs. *Arch. Hydrobiol. Suppl.* 79:69-86
- Strayer, D.L. 1991. Projected distribution of the Zebra Mussel, *Dreissena polymorpha*, in North America. *Canadian Journal of Fisheries and Aquatic Sciences* 48:1389-1395.
- Wu, L. and D.A. Culver. 1991. Zooplankton grazing and phytoplankton abundance: an assessment before and after invasion of *Dreissena polymorpha*. *Journal of Great Lakes Research* 17(4):425-436.
- Yonge, C.M. and J.I. Campbell. 1968. On the heteromyarian condition in the Bivalvia with special reference to *Dreissena polymorpha* and certain Mytilacea. *Transactions of the Royal Society of Edinburgh* 68:21-43.
- Zhadin, V.I. 1952. *Mollusks of fresh and brackish waters of the U.S.S.R.* Academy of Sciences of the U.S.S.R. xvi + 368 pp. [In Russian]. [Israeli Program for Scientific Translations, Jerusalem. 1965.]
- Zhuravel, P.A. 1951. *Dreissena bugensis* (Andrusov) from the Dnieper River System and its recent appearance in the Dnieper Reservoir. *Zoologicheskii Zhurnal* 30:186-188.

#### 4. UNIONOIDEA (Unionidae and Margaritiferidae)

##### a. Handbooks, field guides, regional faunas

- Anderson, R.M., J.B. Layzer and M.E. Gordon. 1991. Recent catastrophic decline of mussels (Bivalvia: Unionidae) in the Little South Fork Cumberland River, Kentucky. *Brimleyana* (17): 1-8.
- Baker, F.C. 1898. The Mollusca of the Chicago area. The Pelecypoda. *The Chicago Academy of Sciences. Bulletin of the Natural History Survey* No. 3(1):1-103, 27 plates.
- Baker, F.C. 1928. The fresh-water Mollusca of Wisconsin. Part II. Pelecypoda. *Bulletin of the Wisconsin Geological and Natural History Survey* 70(2):i-vi + 1-495.
- Bogan, A.E. 1993. *Workshop on Freshwater Bivalves of Pennsylvania*. Sewell, N.J. pp. ii, 1-80, with 11 color plates, 65 figures.
- Bogan, A.E. and P.W. Parmalee. 1983. *Tennessee's Rare Wildlife, Volume II: The Mollusks*. Tennessee Wildlife Resources Agency, Nashville, Tennessee, 123 pp., 53 maps, 53 plates, Volume 2 of Tennessee Wildlife
- Buchanan, A.C. 1980. Mussels (Naiades) of the Meramec River basin, Missouri. *Missouri Department of Conservation, Aquatic Service No. 17*. 76 pp. [4 Figs., 52 Pls., 7 Tabs.]
- Burch, J.B. 1973. Freshwater Unionacean Clams (Mollusca: Pelecypoda) of North America. *Biota of Freshwater Ecosystems Identification Manual* No. 11, U.S. Environmental Protection Agency. + 176 pp.
- Burch, J.B. 1975. *Freshwater Unionacean Clams (Mollusca: Pelecypoda) of North America*, Revised Ed. Malacological Publications (Hamburg, Michigan). 204 pp.



- Burch, J.B. and C.M. Patterson. 1976. Key to the genera of freshwater pelecypods (mussels and clams) of Michigan. [Revised]. *University of Michigan Museum of Zoology, Circular Series* [Revised], No. 4. 37 pp.
- Call, R.E. 1885. A geographic catalogue of the Unionidae of the Mississippi Valley. *Bulletin of the Des Moines Academy of Science* 1(1):5-57.
- Call, R.E. 1895. A study of the Unionidae of Arkansas, with incidental reference to their distribution in the Mississippi Valley. *Transactions of the Academy of Science of St. Louis* 7(1):1-65. 21 plates.
- Call, R.E. 1900. A descriptive catalogue of the Mollusca of Indiana. *Twenty-fourth Annual Report of the Indiana Department of Geology and Natural Resources*, 1899 24:335-535, 1013-1017. Pl. 1-78, figs. 1-13, tables.
- Clarke, A.H. 1973. The freshwater molluscs of the Canadian Interior Basin. *Malacologia* 13(1-2):1-509.
- Clarke, A.H. 1981. The tribe Alasmidontini Unionidae: Anodontinae, Part I, *Pegias*, *Alasmidonta*, and *Arcidens*. *Smithsonian Contributions to Zoology*, No. 326. iii + 101 pp.
- Clarke, A.H. 1982. *The Freshwater Mollusks of Canada*. University of Chicago Press. 446 pp.
- Clarke, A.H. 1985. The tribe Alasmidontini (Unionidae: Anodontinae), Part II: *Lasmigona* and *Simpsonaias*. *Smithsonian Contributions to Zoology* (399):1-75.
- Clarke, A.H. and C.O. Berg. 1959. The freshwater mussels of central New York. *Memoir, Cornell University Agricultural Experimental Station* 367. 80 pp.
- Clench, W.J. 1959. Mollusca. IN: *Fresh Water Biology*, 2nd Ed., W. T. Edmondson, Ed. John Wiley & Sons, Inc. (New York). pp. 1117-1160.
- Clench, W.J. and R.D. Turner. 1956. Freshwater mollusks of Alabama, Georgia, and Florida from Escambia to the Suwannee River. *Bulletin of the Florida State Museum, Biological Sciences* 1(3):97-239. Pls. 1-9, 1 text fig.
- Cummings, K.S. and C.A. Mayer. 1992. Field guide to freshwater mussels of the Midwest. *Illinois Natural History Survey Manual* 5. 194 pp, 78 color figures, 78 maps.
- Frierson, L.S. 1927. *A Classified and Annotated Check List of the North American Naiades*. Baylor University Press (Waco, Texas). 111 pp.
- Goodrich, C. and H. van der Schalie. 1944. A revision of the Mollusca of Indiana. *American Midland Naturalist* 32(2):257-326. Tab. 1.
- Gordon, M.E. and J.B. Lazzer. 1989. Mussels (Bivalvia: Unionoidea) of the Cumberland River. Review of life histories and ecological relationships. *U.S. Dept. of Interior, Fish and Wildlife Service Biological Report* 89(15):i-vii, 1-99.
- Harris, J.L. and M.E. Gordon. 1990. *Arkansas mussels*. Arkansas Game and Fish Commission, Little Rock, Arkansas. 32 pp.
- Heard, W.H. and J.B. Burch. 1966. Key to the genera of freshwater pelecypods (mussels and clams) of Michigan. *Museum of Zoology, University of Michigan Circular* No. 4. 14 pp. Figs. 1-49.
- Johnson, R.I. 1970. The systematics and zoogeography of the Unionidae (Mollusca: Bivalvia) of the southern Atlantic Slope region. *Bulletin of the Museum of Comparative Zoology* 140(6):263-449. Pls. 1-22, Text Figs. 1-5, Tabs. 1-40.
- Johnson, R.I. 1972. The Unionidae (Mollusca: Bivalvia) of peninsular Florida. *Bulletin of the Florida State Museum, Biological Sciences* 16:181-249.
- La Rocque, A. 1967. Pleistocene Mollusca of Ohio. *Bulletin of the Geological Survey of Ohio* 62: vii-xiv, 113-356. Pls. 1-8, Figs. 3-208.
- Lea, I. 1870. *A synopsis of the family Unionidae*, 4th Edition (Philadelphia). 184 pp.
- Mathiak, H.A. 1979. *A River Survey of the Unionid Mussels of Wisconsin 1973 - 1977*. Sand Shell Press (Horicon, Wisconsin). 75 pp.
- Murray, H.D. and A.B. Leonard. 1962. Handbook of Unionid Mussels in Kansas. *University of Kansas Museum of Natural History, Miscellaneous Publications* No. 28. 184 pp. Pls. 1-45, Figs. 1-42, Tabs. 1-5.
- Nalepa, T.F., B.A. Manny, J.C. Roth, S.C. Mozley and D.W. Schloesser. 1991. Long-term decline in freshwater mussels (Bivalvia: Unionidae) of the western basin of Lake Erie. *J. Great Lakes Res.* 17(2): 214-219.
- Oesch, R.D. 1984. *Missouri Naiades: A Guide to the Mussels of Missouri*. Missouri Department of Conservation (Jefferson City). vii + 270 pp.
- Parmalee, P.W. 1967. The fresh-water mussels of Illinois. *Illinois State Museum, Popular Science Series* 8. 108 pp.
- Pennack, R.W. 1979. *Fresh-water Invertebrates of the United States*. Ronald Press Co. (New York).
- Robertson, I.C.S. and C.L. Blakelee. 1948. The Mollusca of the Niagara Frontier region. *Bulletin of the Buffalo Society of Natural Sciences* 19(3):xi + 191 pp.
- Scammon, R.E. 1906. The Unionidae of Kansas, Part 1. *University of Kansas Science Bulletin* 3(9):279-373. Pls. 62-85.
- Sepkoski, J.J., Jr. and M.A. Rex. 1974. Distribution of freshwater mussels: coastal rivers as biogeographic islands. *Systematic Zoology* 23:165-188.
- Smith, D.G. 1991. *Keys to the freshwater macroinvertebrates of Massachusetts*. Dept. of Zoology, University of Massachusetts, Amherst. 256 pp.
- Starnes, L.B. and A.E. Bogan. 1988. The mussels (Mollusca: Bivalvia) of Tennessee. *American Malacological Bulletin*, 6(1):19-37.
- Starrett, W.C. 1971. A survey of the mussels (Unionacea) of the Illinois River: a polluted stream. *Illinois Natural History Survey Bulletin* 30(5):267-403. Col. Pl. 1-4, figs. 1-17, tabs. 1-30.

- Strayer, D.L. 1990. Freshwater Mollusca. pp.335-372. In: B.L. Peckarsky, P.R. Fraissinet, M.A. Penton and D.J. Conklin, jr. *Freshwater macroinvertebrates of northeastern United States*. Comstock Publishing Associates, Ithaca, N.Y.
- Strecker, J.K. 1931. *The distribution of the naiades, or pearly freshwater mussels, of Texas*. Baylor University Museum Special Bulletin 2:1-71.
- Utterback, W.I. 1915. The naiads of Missouri. *American Midland Naturalist* 4:41-53; 69-152; 182-204; 244-273; 311-327; 339-354; 387-400; 432-464.
- Valentine, B.D. and D.H. Stansbery. 1971. An introduction to the naiads of the Lake Texoma region, with notes on the Red River fauna (Mollusca: Unionidae). *Sterkiana* 42:1-40.
- van der Schalie, H. 1938. The naiades (freshwater mussels) of the Cahaba River in Northern Alabama. *Occasional Papers from the Museum of Zoology, University of Michigan* No. 392. 29 pp.
- van der Schalie, H. 1940. The naiad fauna of the Chipola River, in northern Florida. *Lloydia* 3:191-208.
- van der Schalie, H. and A. van der Schalie. 1950. The mussels of the Mississippi River. *American Midland Naturalist* 44(2):448-466.
- Vidrine, M.F. 1993. *The historical distributions of freshwater mussels in Louisiana*. Gali. Q. vidrine Collectables. Eunice, Louisiana. pp. xii, 1-225, 138 maps, 20 color plates.
- Walker, B. 1918. A synopsis of the classification of fresh-water Mollusca of North America, north of Mexico, and a catalogue of the more recently described species, with notes. *Miscellaneous Publications of the Museum of Zoology, University of Michigan* No. 6:1-213.
- Walker, B. 1918. Notes on North American naiades, I. *Occasional Papers of the Museum of Zoology, University of Michigan* No. 49:1-6.
- Walker, B. 1918. The Unionia fauna of the Great Lakes. *Miscellaneous Publications of the Museum of Zoology, University of Michigan*. 40:1-83.
- Walker, B. and B.H. Wright. 1902. *Checklist of North American Naiades*. Detroit, printed for the authors, 19 pp.
- Watters, G.T. 1992. Unionids, fishes, and the species-area curve. *Journal of Biogeography* 19:481-490.
- Watters, G.T. 1993. *A guide to the freshwater mussels of Ohio*. Revised edition. The Division of Wildlife, The Ohio Department of Natural Resources. 106 pp.
- Williams, J.C. and G.A. Schuster. 1989. *Freshwater mussel investigations of the Ohio River Mile 317.0 to Mile 381.0*. Kentucky Department of Fish and Wildlife Resources, Division of Fisheries. pp. iii, 1-57, 8 colored plates.
- Williams, J.D., S.L.H. Fuller, and R. Grace. 1992. Effects of impoundment on freshwater mussels (Mollusca: Bivalvia: Unionidae) in the main channel of the Black Warrior and Tombigbee rivers in western Alabama. *Bulletin of the Alabama Museum of Natural History*. 13:1-10.
- Williams, J.E. and R.J. Neves. 1992. Introducing the elements of biological diversity in the aquatic environment. In J.E. Williams and R.J. Neves (eds.), *Biological diversity in aquatic management. Transactions of the 57th North American Wildlife and Natural Resources Conference*. (1992):345-354

#### b. Taxonomy of unionids

- Bereza, D.J. and S.L.H. Fuller. 1975. Notes on "*Lampsilis* " *ochracea* (Say) (Mollusca: Bivalvia). *ASB Bulletin* 22:42.
- Bereza, D.J., M.F. Vidrine and S.L.H. Fuller. 1976. Anatomical differences between *Ligumia nasuta* (Say) and *L. subrostrata* (Say) (Mollusca: Bivalvia: Unionidae). *ASB Bulletin* 23(2):43.
- Binney, W.G. and G.W. Tryon, Jr. 1864. *The Complete Writings of Constantine Smltz Rafinesque on Recent and Fossil Conchology*. Bailliere Brothers (New York). 103 pp.
- Bogan, A.E. 1985. A comment on Strophitinae Gordon, 1981 (Unionidae, Bivalvia). *The Nautilus*, 99(4):141-142.
- Bogan, A.E. 1988. A bibliographic history of C.S. Rafinesque's work on North American freshwater bivalves. *Archives of Natural History* 15(2):149-154.
- Clarke, A.H., Jr. and W.J. Clench. 1965. *Amblema* Rafinesque, 1820 (Lamellibranchiata): proposed addition to the official list and proposed suppression of *Amblema* Rafinesque, 1819. Z.N.(S.) 1699. *Bulletin of Zoological Nomenclature* 22(3):196-197.
- Clarke, A.H. and W.J. Clench. 1966. Revision of proposals concerning *Amblema* Rafinesque, 1820. *Bulletin of Zoological Nomenclature* 22(5/6):341.
- Davis, G.M. 1983. Relative roles of molecular genetics, anatomy, morphometrics and ecology in assessing relationships among North American Unionidae (Bivalvia). *Systematics Association Special Volume* No. 24, Protein Polymorphism: Adaptive and Taxonomic Significance, G. S. Oxford and D. Rollinson, Eds. Academic Press (London). pp. 193-222.
- Davis, G.M. 1984. Genetic relationships among some North American Unionidae (Bivalvia): sibling species, convergence, and cladistic relationships. *Malacologia* 25(2):629-648.
- Davis, G.M. and S.L.H. Fuller. 1981. Genetic relationships among recent Unionacea (Bivalvia) of North America. *Malacologia* 20(2):217-253. 4 Figs., 14 Tabs., 2 Appendices.
- Davis, G.M., W.H. Heard, S.L.H. Fuller and C. Hesterman. 1981. Molluscan genetics and speciation in *Elliptio* and its relationship to other taxa of North American Unionidae (Bivalvia). *Biological Journal of the Linnean Society* 15:131-150.

- Haas, F. 1969. Superfamily Unionacea. IN: *Treatise on Invertebrate Paleontology*, R. C. Moore, Ed. Geological Society of America (Boulder, Colorado). Part N, Volume 1 (of 3), Mollusca 6, Bivalvia. pp. N411-N470.
- Haas, F. 1969. Superfamilia Unionacea. *Das Tierreich* (Berlin). x + 663 pp.
- Heard, W.H. and R.H. Gukert. 1970. A re-evaluation of the recent Unionacea (Pelecypoda) of North America. *Malacologia* 10(2):333-355.
- Hoeh, W.R. 1990. Phylogenetic relationships among eastern North American *Anodonta* (Bivalvia: Unionidae). *Malacological Review* 23:63-82.
- Hoeh, W.R. 1991. The evolution and consequences of simultaneous hermaphroditism in the freshwater mussel genus *Utterbackia* (Bivalvia: Unionidae). Unpublished Ph.D. Dissertation, University of Michigan, Ann Arbor. 97 pp.
- Hoeh, W.R. and J.B. Burch. 1989. The taxonomic status of *Anodonta lacustris* Lea (Bivalvia: Unionidae). *Walkerana* 3(10):263-276.
- Johnson, R.I. 1978. Systematics and zoogeography of *Plagiola* (= *Dysnomia* = *Epioblasma*), an almost extinct genus of freshwater mussels (Bivalvia: Unionidae) from middle North America. *Bulletin of the Museum of Comparative Zoology* 148:239-320.
- Kat, P.W. 1983. Genetics, morphological divergence, and speciation among *Lampsilis*. *Journal of Molluscan Studies* 49:133-145.
- Kat, P.W. 1983. Genetics and morphological divergence among nominal species of North American *Anodonta* (Bivalvia: Unionidae). *Malacologia* 23(2):361-374.
- Ortmann, A.E. 1912. Notes upon the families and genera of the naiades. *Annals of the Carnegie Museum* 8(2):222-365. Pls. 18-20, Figs. 1-28.
- Ortmann, A.E. and B. Walker. 1922. On the nomenclature of certain North American naiades. *Occasional Papers of the Museum of Zoology, University of Michigan* No. 112. 75 pp.
- Simpson, C.T. 1900. Synopsis of the naiades, or pearly freshwater mussels. *Proceedings of the U.S. National Museum* 22(1205):501-1044.
- Simpson, C.T. 1914. *A descriptive catalogue of the naiades or pearly freshwater mussels*. Bryant Walker (Detroit). Parts I-III. xi + 1540 pp.
- Stansbery, D.H. 1970. Comments on *Pleurobema coccineum* (Conrad, 1836) in the Muskingham River System. *Pittsburgh Shell Club Bulletin*. No. 5:15.
- Stansbery, D.H. 1983. Some sources of nomenclatural and systematic problems in unionid mollusks. IN: *Report of Freshwater Mussels Workshop*; 26-27 October 1982, A. C. Miller, Ed. U.S. Army Engineer Waterways Experiment Station (Vicksburg, MS). pp. 46-62.
- Turgeon, D.D., A.E. Bogan, E.V. Coan, W.K. Emerson, W.G. Lyons, W.L. Pratt, C.F.E. Roper, A. Scheltema, F.G. Thompson, and J.D. Williams. 1988. Common and scientific names of aquatic invertebrates from the United States and Canada: Mollusks. *American Fisheries Society Special Publication* 16:vii, 1-277, 12 color plates.
- van der Schalie, H. 1951. An old problem in naiad nomenclature. *Bulletin of the American Malacological Union* 1951:4-5. (Reprinted in *The Nautilus* 65(3):93-99. 1952; *Sterkiana* 9:30-31. 1963.)
- Vokes, H.E. 1980. *Genera of the Bivalvia: A Systematic and Bibliographic Catalogue*. Paleontological Research Institution (Ithaca, New York).

### c. Literature on Maryland unionids

- Anonymous. 1994. Maryland DNR declares four species extinct. *Ecology USA* 23(20):196.
- Bartgis, R. and L.H. MacIvor. 1993. Brook floater surveys in Maryland. Report to Chesapeake Bay and Endangered Species Fund and the U.S. Fish and Wildlife Service. Study No. E-3-6. Final report. [5 pp], 4 tables.
- Cohen, R.R.H., P.V. Dresler, E.J.P. Phillips, and R.L. Cory. 1984. The effect of the Asiatic clam, *Corbicula fluminea*, on phytoplankton of the Potomac River, Maryland. *Limnology and Oceanography* 29(1):170-180.
- Counts, C.L., III, and T.L. Bashore. 1991. Mollusca of Assateague Island, Maryland and Virginia: a reexamination after seventy-five years. *The Veliger* 34(2):214-221.
- Counts, C.L., III, T.S. Handwerker, and R.V. Jesien. 1991. The naiades (Bivalvia: Unionoidea) of the Delmarva Peninsula. *American Malacological Bulletin* 9(1):27-37.
- Flynn K.C. & W.T. Mason (eds.). 1978. *The Freshwater Potomac: Aquatic communities and environmental stresses*. Interstate Commission on the Potomac River Basin. Rockville MD. 194 pp.
- Fuller, S.L.H. 1978. Changes in the molluscan community of the Middle Potomac River during the past two decades. pp. 124-131 in K.C. Flynn & W.T. Mason (eds.). *The Freshwater Potomac: Aquatic communities and environmental stresses*. Interstate Commission on the Potomac River Basin, Rockville, MD., 194 pp.
- Gerberich, A.G. 1984. *The endangered and threatened freshwater mollusks of Maryland*. Maryland Natural Heritage Program Special Publication, 84-1:245-266.
- Girard, C. 1856. Catalogue of Recent shells and other molluscs found in the District of Columbia, prepared from specimens in the cabinet of Dr. E. Foreman. *Proceedings of the National Institute, Washington, D.C. N.S.* 1(2):78-82.
- Kat, P.W. 1982. Effects of population density and substratum type on growth and migration of *Elliptio complanata* (Bivalvia: Unionidae). *Malacological Review* 15:119-127.



- Lehnert, C. 1885. A list of Recent land and fresh-water mollusks of the District of Columbia and vicinity. *Pastime* 3(8):5-8.
- Long, G.A. 1979. Anodonta and Lampsilis at Loch Raven Reservoir. (Abstract). *Bulletin of the American Malacological Union, Inc.* 1979:67.
- Long, G.A. 1983. The unionids (Bivalvia) of Loch Raven Reservoir, Maryland. *The Nautilus* 97(3):114-116.
- Marshall, W.A. 1917. Lampsilis ventricosa cohongoronta in the Potomac River. *The Nautilus* 31(2):40-41.
- Marshall, W.B. 1918. Lampsilis ventricosa cohongoronta in the Potomac River valley. *The Nautilus* 32(2):51-53.
- Marshall, W.B. 1930. Lampsilis ventricosa cohongoronta in the Potomac River. *The Nautilus* 44(1):19-21.
- Marshall, W.B. 1930. Mollusks from below Conowingo Dam, Maryland. *The Nautilus* 43:87-88.
- Maryland Department of Natural Resources. 1992. *Rare, threatened and endangered animals of Maryland*. Maryland Department of Natural Resources, Annapolis, MD. 9 pp.
- Morrison, J.P.E. 1975. Maryland and Virginia mussels of Lister. *Bulletin of the American Malacological Union, Inc.* 1974:36-39.
- Neves, R.J. 1983. The status of freshwater mussel research in Virginia. pp. 155-168 in A.C. Miller (compiler) *Report of freshwater mollusks workshop (26-27 October 1982)*. U.S. Army Engineer Waterways Experimental Station Vicksburg Mississippi. 196 pp.
- Neves, R.J. 1991. Mollusks. pp. 251-320 in K. Terwilliger (ed.). *Virginia's Endangered Species. Proceedings of a Symposium*. Department of Game and Inland Fisheries Commonwealth of Virginia. 672 pp.
- Neves, R.J., G.B. Pardue, E.F. Benfield, and S.D. Dennis. 1980. An evaluation of endangered mollusks in Virginia. Virginia Commission on Game and Inland Fisheries Project Report E-F- 1. 140 pp.
- Ortmann, A.E. 1912. Lampsilis ventricosa (Barnes) in the Upper Potomac drainage. *The Nautilus* 26(4):51-54.
- Ortmann, A.E. 1913. the Alleghenian Divide, and its influence on the freshwater fauna. *Proceedings of the American Philosophical Society* 52(210):287-396, pls. 12-14. ???
- Ortmann, A. E. 1919. A monograph on the naiades of Pennsylvania. Part 3. Systematic account of the genera and species. *Memoirs of the Carnegie Museum* 8(1):xiv + 384 pp. 21 Pls., 34 Figs.
- Ortmann, A. E. 1920. Correlation of shape and station in freshwater mussels. *Proceedings of the American Philosophical Society* 59(4):269-312. 1 Map.
- Pilsbry, H.A. 1894. Critical list of mollusks collected in the Potomac Valley. *Proceedings of the Academy of Natural Sciences of Philadelphia* 46:11-31, pl. 1.
- Reardon, L. 1929. a contribution to our knowledge of the anatomy of the fresh-water mussels of the District of Columbia. *Proceedings of the United States National Museum* 75:1-12, pls. 1-5.
- Richards, H.G. 1934. A list of the mollusks of the District of Columbia and vicinity. *American Midland Naturalist* 15(1):85-88.
- Stansbery, D.H. 1987. Identification of shell remains from prehistoric site 46HM73 on the upper Potomac River in West Virginia. *West Virginia Archaeologist* 39(2):57-58.
- Taylor, R.W. 1985. Comments on the distribution of freshwater mussels (Unionacea) of the Potomac River headwaters in West Virginia. *The Nautilus* 99(2-3):84-87.
- Tryon, G.W. 1861. On the Mollusca of Harper's Ferry Virginia. *Proceedings of the Academy of Natural Sciences of Philadelphia* 13:396-399.

#### d. Endangered species

- Anonymous. 1980. The ten most endangered species in North America. *North Dakota Outdoors* 42(12):20-21.
- Athearn, H.D. 1970. Discussion of Dr. Heard's paper. Symposium on Endangered Mollusks. *Malacologia* 10(1):28-31.
- Bickel, D. 1973. Critique "Proceedings of a symposium on rare and endangered mollusks (Naiads) of the U.S. Columbus, Ohio 1971". *Sterkiana*. 49:18-20.
- Bogan, A.E. 1993. Freshwater bivalve extinctions: search for a cause. *American Zoologist*, 33:599-609.
- Bogan, A.E., and P.W. Parmalee. 1983. *Tennessee's rare wildlife. Volume II: The Mollusks*. Tennessee Wildlife Resources Agency and Tennessee Department of Conservation. 123 pp.
- Branson, B.A., D.F. Harker, Jr., J.M. Basin, M.E. Medley, D.L. Batch, M.L. Warren, Jr., W.H. Davis, W.C. Houtcooper, B. Monroe, Jr., L.R. Phillippe, and P. Cupp. 1981. Endangered, threatened, and rare animals and plants of Kentucky. *Transactions of the Kentucky Academy of Science*. 42(3-4):77-89.
- Case, D.S. 1986. Endangered wildlife management in Ohio. in S.K. Majumdar, F.J. Brenner, and A.F. Rhodes (eds.). *Endangered and threatened species programs in Pennsylvania and other states: Causes, Issues and Management*. *Pennsylvania Academy of Science*. 388-394.
- Chambers, S.M. 1981. Protection of mollusks under the Endangered Species Act of 1973. *Bulletin of the American Malacological Union, Inc.* 1981:55-59.
- Clark, C.F. 1971. Management of naiad populations in Ohio. in S.E. Jorgenson and R.E. Sharp (eds.). *Proceedings of a symposium on rare and endangered mollusks (naiads) of the U.S. Region 3*, Bureau Sport Fisheries and Wildlife, U.S. Fish Wildlife Service. Twin Cities, Minnesota. pp 26-33.
- Dundee, D.S. 1981. Endangered species - why worry? *Bulletin of the American Malacological Union, Inc.* 1981:54-55.

- Fikes, M.A., and R.A. Tubb. 1971. *Amblema plicata* as a pesticide monitor. in S.E. Jorgenson and R.E. Sharp (eds.). *Proceedings of a symposium on rare and endangered mollusks (naiads) of the U.S. Region 3*, Bureau Sport Fisheries and Wildlife, U.S. Fish Wildlife Service. Twin Cities, Minnesota. pp 34-37.
- Fuller, S.L.H. 1977. Freshwater and terrestrial mollusks. pp. 143-194 in J.E. Cooper, S.S. Robinson, and J.B. Funderburg (eds.). *Endangered and threatened plants and animals of North Carolina*. North Carolina State Museum of Natural History, Raleigh.
- Fuller, S.L.H. 1980. Fresh-water mussels (Mollusca: Bivalvia: Unionidae). pp. 114-125. In: *Proceedings of the first South Carolina Endangered Species Symposium*. D.M. Forsythe and W.B. Ezell (eds.) South Carolina Wildlife and Marine Resources Dept., Columbia. 201 pp.
- Gerberich, A.G. 1984. The endangered and threatened freshwater mollusks of Maryland. pp. 245-266. In: *Threatened and endangered plants and animals of Maryland*. A.W. Norden, D.C. Forester, and G.H. Fenwick (eds.). *Maryland Natural Heritage Program Special Publication 84-1*:1-475.
- Harris, J.L., and M.E. Gordon. 1988. Distribution and status of rare and endangered mussels (Mollusca: Margaritiferidae, Unionidae) in Arkansas. *Arkansas Academy of Science Proceedings*. 41(1987):49-56.
- Havlik, M.E. 1980. The historic and present distribution of the endangered naiad mollusk *Lampsilis higginsii* (Lea, 1857). *Bulletin of the American Malacological Union, Inc.* 1980:19-22.
- Heard, W. H. 1970. 3. Eastern freshwater mollusks. (II). The south Atlantic and Gulf drainages. IN: *Papers on the Rare and Endangered Mollusks of North America*, A. H. Clarke, Ed. *Malacologia* 10:1-56.
- Holt, F. T. 1974. *Rare and endangered species of Missouri*. Missouri Department of Conservation and U.S.D.A., S.C.S. unpaginated.
- Imlay, M.J. 1972. Our rivers... Death beds for endangered mollusks. *Minnesota Volunteer*. 35:30-39.
- Jenkinson, J.J. 1981. Endangered or threatened aquatic mollusks of the Tennessee River System. *Bulletin of the American Malacological Union, Inc.* 1981:43-45.
- Knight, K.B. 1986. Threatened and endangered species in West Virginia. in S.K. Majumdar, F.J. Brenner, and A.F. Rhodes (eds.). *Endangered and threatened species programs in Pennsylvania and other states: Causes, Issues and Management*. *Pennsylvania Academy of Science*. 410-427.
- Lowe, D.W., J.R. Matthews, and C.J. Moseley (eds.). 1990. *The official World Wildlife Fund guide to endangered species of North America*. Beacham Publishing, Inc. Washington D.C. 2:954-1021.
- Michigan Department of Natural Resources. 1976. *Michigan's endangered and threatened species program*. Michigan Department of Natural Resources (Lansing). 30 pp.
- Miller, A.C. 1981. Endangered mollusk studies at the U.S. Army Engineer Waterways Experiment Station Vicksburg, Mississippi. *Bulletin of the American Malacological Union, Inc.* 1981:45-49.
- Miller, A.C., B.S. Payne, and T. Siemsen. 1986. Description of the habitat of the endangered mussel *Plethobasus cooperianus*. *The Nautilus*. 100(1):14-18.
- Neves, J.J. 1991. Mollusks. pp 251-320. IN: K. Terwilliger (ed.) *Virginia's Endangered Species: proceedings of a symposium*. the McDonald and Woodward publishing Company, Blacksburg, Virginia.
- Neves, R.J., G.B. Pardue, E.F. Benfield, and S.D. Dennis. 1980. An evaluation of endangered mollusks in Virginia. Virginia Commission on Game and Inland Fisheries Project Report E-F- 1. 140 pp.
- Nordstrom, G.R., W.L. Pflieger, K.C. Sadler, and W.H. Lewis. 1977. *Rare and endangered species of Missouri*. Missouri Department of Conservation, U.S. Department of Agriculture and the Soil Conservation Service. pp. 129.
- Randolph, J.P. 1986. Virginia's endangered species program. in S.K. Majumdar, F.J. Brenner, and A.F. Rhodes (eds.). *Endangered and threatened species programs in Pennsylvania and other states: Causes, Issues and Management*. *Pennsylvania Academy of Science*. 395-409.
- Ruhr, G. 1971. The need and basis for action programs for endangered species. in S.E. Jorgenson and R.E. Sharp (eds.). *Proceedings of a symposium on rare and endangered mollusks (naiads) of the U.S. Region 3*, Bureau Sport Fisheries and Wildlife, U.S. Fish Wildlife Service. Twin Cities, Minnesota. pp 42-46.
- Shiffer, C. 1985. Drainage patterns in Pennsylvania. In: H.H. Genoways and Fred J. Brenner (eds.) *Species of special concern in Pennsylvania*. *Special Publication of Carnegie Museum of Natural History* No. 11
- Stansbery, D.H. 1970. Eastern freshwater mollusks. (I.) The Mississippi and St. Lawrence River systems. American Malacological Union Symposium on Rare and Endangered Mollusks. *Malacologia* 10(1): 9-22.
- Stansbery, D.H. 1971. Rare and endangered mollusks in the Eastern United States. in S.E. Jorgenson and R.E. Sharp (eds.). *Proceedings of a symposium on rare and endangered mollusks (naiads) of the U.S. Region 3*, Bureau Sport Fisheries and Wildlife, U.S. Fish Wildlife Service. Twin Cities, Minnesota. pp 5-18f, 50 fig.
- Stansbery, D.H. 1976. Naiad mollusks. *Endangered and Threatened Plants and Animals of Alabama*. *Bulletin of the Alabama Museum of Natural History*. 2:42-52.
- Stein, C.B. 1971. Naiad life cycles: Their significance in the conservation of the fauna. in S.E. Jorgenson and R.E. Sharp (eds.). *Proceedings of a symposium on rare and endangered mollusks (naiads) of the U.S. Region 3*, Bureau Sport Fisheries and Wildlife, U.S. Fish Wildlife Service. Twin Cities, Minnesota. pp 19-25 + 3 plates.
- Stein, C.B., and M.J. Imlay. 1976. Current trends in Malacology: Environmental impact studies and endangered species research. *Bulletin of the American Malacological Union, Inc.* 1976:62-67.
- Sterns, E. and D. Felder. 1978. Identification of host fishes for four species of freshwater mussels. *American Midland Naturalist* 100:233-236.

- U.S. Department of Interior, Fish and Wildlife Service (USDI/FWS). 1994. Endangered and threatened wildlife and plants; Animal candidate review for listing as endangered or threatened species. 50 Code of Federal Regulations Part 17, *Federal Register*. 59(219):58982-59028.
- U.S. Department of Interior, Fish and Wildlife Service. 1993. Endangered & threatened wildlife and plants. 50 Code of Federal Regulations 17.11 & 17.12. Part 17. pp. 1-40
- van der Schalie, H. 1975. An ecological approach to rare and endangered species in the Great Lakes region. *The Michigan Academician*. 8:7-22.
- Warren, M.L., Jr., W.H. Davis, R.R. Hannan, M. Evans, D.L. Batch, B.D. Anderson, B. Palmer-Ball, Jr., J.R. MacGregor, R.R. Cicerello, R. Athey, B.A. Branson, G.L. Fallo, B.M. Burr, M.E. Medley, and J.M. Baskin. 1986. Endangered, threatened, and rare plants and animals of Kentucky. *Transactions of the Kentucky Academy of Science*. 47(3-4): 83-98.
- Wilson, J.H. 1984. (ed.) *Rare and endangered species of Missouri*. Missouri Department of Conservation. Conservation Commission of the State of Missouri.

#### e. Glochidia, reproduction and fish hosts

- Allen, E. 1924. The existence of a short reproductive cycle in *Anodonta imbecilis*. *Biological Bulletin* 46:88-94.
- Arey, L.B. 1921. An experimental study on glochidia and the factors underlying excystment. *Journal of Experimental Zoology* 33(3):463-499. Pls. 1-3.
- Arey, L.B. 1923. Observations on an acquired immunity to a metazoan parasite. *Journal of Experimental Zoology* 38(3):377-381.
- Arey, L.B. 1924. Glochidial cuticulae, teeth, and the mechanics of attachment. *Journal of Morphology and Physiology* 39:323-335.
- Arey, L.B. 1932. The formation & structure of the glochidial cyst. *Biological Bulletin* 62(2):212-221.
- Arey, L.B. 1932. A microscopical study of glochidial immunity. *Journal of Morphology*. 53:367-378.
- Arey, L.B. 1932. The nutrition of glochidia during metamorphosis. *Journal of Morphology*. 53:201-221.
- Bruenderman, S.A. and R.J. Neves. 1993. Life history of the endangered fine-rayed pigtoe *Fusconaia cuneolus* (Bivalvia: Unionidae) in the Clinch River, Virginia. *American Malacological Bulletin*. 10(1):83-91.
- Chamberlain, T.K. 1934. The glochidial conglomerates of the Arkansas fanshell, *Cyprogenia aberti* (Conrad). *Biological Bulletin* 66:55-61.
- Clark, H.W. and S. Stein. 1921. Glochidia in surface towings. *The Nautilus* 35(1):16-20.
- Conner, C.H. 1905. Glochidia of *Unio* on fishes. *The Nautilus* 18(12):142-143.
- Conner, C.H. 1907. The gravid periods of Unios. *The Nautilus* 21(8):87-89.
- Conner, C.H. 1909. Supplementary note on the breeding seasons of the Unionidae. *The Nautilus* 22(10):111-112. 1 Tab.
- Davenport, D. and M. Warmuth. 1965. Notes on the relationship between the freshwater mussel *Anodonta imbecilis* Say and the alewife *Pomolobus pseudoharengus* Wilson. *Limnology and Oceanography* 10:R74-R78.
- Dechtiar, A.O. 1972. New parasite records for Lake Erie fish. *Technical Reports, Great Lakes Fisheries Commission* No. 17. iii + 20 pp.
- D'Eliscu, P.N. 1972. Observation of the glochidium, metamorphosis, and juvenile of *Anodonta californiensis*, Lea, 1857. *The Veliger* 15:57-58.
- Dietz, T.H., H. Silverman, and W.L. Steffens. 1985. Movement of maternal calcium into glochidia in a unionid. *American Zoologist*. 25(4):57a.
- Edgar, A.L. 1965. Observations on the sperm of the pelecypod *Anodontoides ferussacianus* (Lea). *Transactions of the American Microscopical Society* 84:228-230.
- Ellis, M.M. and M.D. Ellis. 1926. Growth and transformation of parasitic glochidia in physiological nutrient solutions. *Science* 54(1667):579-580.
- Ellis, M.M., and M.D. Ellis. 1927. Responses of closed glochidia to fish-blood equivalents of sodium, potassium, and calcium salts. *Journal of Experimental Zoology*. 47:197-223.
- Ellis, M.M. and M. Keim. 1918. Notes on the glochidia of *Strophitus edentulus pavonius* (Lea) from Colorado. *The Nautilus* 32:17-18.
- Fuller, S.L.H. 1974. Clams and mussels. IN: *Pollution Ecology of Freshwater Invertebrates*, C.W. Hart, Jr. and S.L.H. Fuller, Eds. Academic Press (New York). pp. 215-273. [host fishes]
- Fustish, C.A. and R.E. Millemann. 1978. Glochidiosis of salmonid fishes. 2. Comparison of tissue response of coho and chinook salmon to experimental infection with *Margaritifera margaritifera* (L.) (Pelecypoda: Margaritiferidae). *Journal of Parasitology* 64(1):155-157.
- Giusti, F. 1973. The minute shell structure of the glochidium of some species of the genera *Unio*, *Potomida* and *Anodonta* (Bivalvia: Unionacea). *Malacologia* 14(1-2):291-301. 26 Pls.
- Hanek, G. and C.H. Fernando. 1978. Spatial distribution of gill parasites of *Lepomis gibbosus* (L.) and *Ambloplites rupestris* (Raf.). *Canadian Journal of Zoology* 56(6): 1235-1240.
- Hanek, G. and C.H. Fernando. 1978. The role of season, habitat, host age and sex on gill parasites of *Lepomis gibbosus* (L.). *Canadian Journal of Zoology* 56(6):1247-1250.
- Hanek, G. and C.H. Fernando. 1978. The role of season, habitat, host age and sex of gill parasites of *Ambloplites rupestris* (Raf.). *Canadian Journal of Zoology* 56(6):1251-1253.



- Heard, W.H. 1975. Sexuality and other aspects of reproduction in *Anodonta* (Pelecypoda: Unionidae). *Malacologia* 15(1):81-103.
- Heard, W.H. and S.S. Hendrix. 1964. Behavior of unionid glochidia. *Bulletin of the American Malacological Union* 1964:2-4.
- Heffelfinger, K.A. 1969. Studies of the structure and ultrastructure of the glochidial stage of the naiad *Actinonaias ligamentina* (Lamarck, 1819) (Mollusca: Bivalvia: Unionidae). M.S. Thesis, Ohio State University, Columbus, Ohio.
- Hill, A.M. 1987. The glochidia (larvae) of the freshwater mussel *Margaritifera hembeli* (Unionacea: Margaritiferidae) fish host species, morphology and period of fish host infection. M.S. Thesis, Northwestern State University, Louisiana.
- Hoggarth, M.A. 1986. The mechanics of glochidial attachment (Mollusca: Bivalvia: Unionidae). *American Malacological Bulletin*. 4(1):117-118.
- Hoggarth, M.A. 1987. Determination of anterior-posterior orientation of glochidia by the examination of glochidial valves present within the umbos of juvenile unionid clams (Mollusca: Bivalvia). *Ohio Journal of Science*. 87(3):93-95.
- Hoggarth, M.A. 1988. The use of glochidia in the systematics of the Unionidae (Mollusca: Bivalvia). Ph.D. Thesis, Dept. of Zoology, Ohio State University. pp i-xxiv, 1-340.
- Hoggarth, M.A. 1992. An examination of the glochidia-host relationships reported in the literature for North American species of Unionacea (Mollusca: Bivalvia). *Malacology Data Net* 3(1-4):1-30.
- Hoggarth, M.A. 1994. A key to the glochidia of the Unionidae of Ohio. M.A. Hoggarth, Dept. of Life and Earth Sciences, Otterbein College, Westerville, Ohio. [13 pp], 6 figs.
- Hoggarth, M.A. 1994. A key to the glochidia of the Unionidae of Texas. M.A. Hoggarth, Dept. of Life and Earth Sciences, Otterbein College, Westerville, Ohio. [12 pp], 5 figs.
- Hoggarth, M.A., and A.S. Gaunt. 1988. Mechanics of glochidial attachment (Mollusca: Bivalvia: Unionidae). *Journal of Morphology*. 198(1):71-81.
- Howard, A.D. 1914. A second case of metamorphosis without parasitism in the Unionidae. *Science* 40(1027):353-355. 1 Tab.
- Howard, A.D. 1914. Experiments in propagation of fresh-water mussels of the *Quadrula* group. *Report of the U.S. Commissioner of Fisheries for 1913*. Appendix IV. pp. 1-52. Pl. I-VI, Tabs. 1-4. [Issued separately as Bureau of Fisheries Document 801].
- Howard, A.D. 1915. Some experimental cases of breeding among the Unionidae. *The Nautilus* 29:4-11.
- Howard, A.D. 1917. An artificial infection with glochidia on the river herring. *Transactions of the American Fisheries Society*. 46(2):93-100.
- Howard, A.D. 1922. Experiments in the culture of fresh-water mussels. *Bulletin of the U.S. Bureau of Fisheries* 38:1-89. [Also issued as U.S. Bureau of Fisheries Document No. 916]
- Howard, A.D. 1951. A river mussel on a salamander. *Natural History Miscellanea*. #77:1-6.
- Howard, A.D. and B.J. Anson. 1922. Phases in the parasitism of the Unionidae. *Journal of Parasitology* 9:68-82.
- Hudson, R.G. and B.G. Isom. 1984. Rearing juveniles of the freshwater mussels (Unionidae) in a laboratory setting. *The Nautilus* 98(4):129-135.
- Ingram, W.M. 1941. Tadpoles as hosts for the glochidia of the freshwater mussel. *Current Science*. 10:535-536.
- Isom, B.G. 1983. Potential uses of in vitro culture of freshwater mussel glochidia for conservation. IN: *Report of Freshwater Mussel Workshop*; 26-27 October 1982, A.C. Miller, Ed. U.S. Army Engineer Waterways Experiment Station (Vicksburg, Mississippi). pp. 42-45.
- Isom, B.G. and R.G. Hudson. 1982. In vitro culture of parasitic freshwater mussel glochidia. *The Nautilus* 96(4):147-151.
- Isom, B.G., and R.G. Hudson. 1984. Freshwater mussels and their fish hosts: physiological aspects. *Journal of Parasitology*. 70(2):318-319.
- Kakonge, S.A.K. 1972. The ecology of some metazoan parasites of, and their effect on, some stream fishes and fry. Ph.D. Dissertation, University of Waterloo. 163 pp.
- Karna, D.W. and R.E. Millmann. 1978. Glochidiosis of salmonid fishes. 3. Comparative susceptibility to natural infection with *Margaritifera margaritifera* (L.) (Pelecypoda: Margaritiferidae) and associated histopathology. *Journal of Parasitology* 64(3):528-537.
- Kat, P.W. 1984. Parasitism and the Unionacea (Bivalvia). *Biological Review* 59:189-207.
- Kozel, T.R. and F.H. Whittaker. 1982. Ectoparasite of the rainbow darter *Etheostoma caeruleum* Storer, from Harrods Creek, Oldham County, Kentucky. *Proceedings of the Helminthological Society of Washington* 49(1):138-139.
- Lefevre, G. and W.C. Curtis. 1908. Experiments in the artificial propagation of fresh-water mussels. *Bulletin of the Bureau of Fisheries* 28:615-626.
- Lefevre, G. and W.C. Curtis. 1910. Reproduction and parasitism in the Unionidae. *Journal of Experimental Zoology* 9(1):79-115. Pls. 1-5, Fig. 1.
- Lefevre, G. and W.C. Curtis. 1911. Metamorphosis without parasitism in Unionidae. *Science* 33(857): 863-865.
- Lefevre, G. and W.C. Curtis. 1912. Studies on the reproduction and artificial propagation of freshwater mussels. *Bulletin of the Bureau of Fisheries* 30:103-201. Pls. 6-17, Figs. 1-4, 1 Tab. [Also published as U.S. Bureau of Fisheries Document 756. Reprinted in *Sterkiana* 47, 48 (1972), 49, 51 (1973), 57 (1975), 61, 63-64 (1976)]

- Leidy, J. 1883. On the reproduction of parasites of *Anodonta fluviatilis*. *Proceedings of the Academy of Natural Sciences, Philadelphia*. 35:44-46.
- Lillie, F.R. 1893. Account of the embryology of *Unio complanata*. *Journal of Morphology* 8:569-578. 1 Pl.
- Lillie, F.R. 1895. The embryology of the Unionidae. *Journal of Morphology* 10:1-100. 6 Pls.
- Margolis, L. 1982. Parasitology of Pacific salmon - an overview. IN: *Aspects of Parasitology. A Festschrift Dedicated to the Fiftieth Anniversary of the Institute of McGill University, 1932-1982*, E. Meerovitch, Ed. McGill University (Montreal). pp. 135-226.
- Margolis, L. and J.R. Arthur. 1979. Synopsis of the parasites of fishes of Canada. *Bulletin of the Fisheries Research Board of Canada* 199:vi + 269 pp.
- Merrick, A.D. 1930. Some quantitative determinations of glochidia. *The Nautilus*. 43(3):89-91.
- Meyers, T.R., R.E. Millemann and C.A. Fustish. 1980. Glochidiosis of salmonid fishes. 4. Humoral and tissue responses of coho and chinook salmon to experimental infection with *Margaritifera margaritifera* (L.) (Pelecypoda: Margaritiferidae). *Journal of Parasitology* 66(2):274-281. Illustrated.
- Neves, R.J., L.R. Weaver, and A.V. Zale. 1985. An evaluation of host fish suitability for glochidia of *Villosa vanuxemi* and *V. nebulosa* (Pelecypoda: Unionidae). *American Midland Naturalist*. 113(1):13-19.
- Neves, R.J., and J.C. Widlak. 1988. Occurrence of glochidia in stream drift and on fishes of the Upper North Fork Holston River, Virginia. *American Midland Naturalist*. 119(1):111-120.
- Ortmann, A.E. 1909. The breeding season of Unionidae in Pennsylvania. *The Nautilus* 22:91-95.
- Ortmann, A.E. 1909. The breeding season of Unionidae in Pennsylvania. *The Nautilus* 22: 99-103.
- Ortmann, A.E. 1910. The discharge of the glochidia in the Unionidae. *The Nautilus* 24:94-95.
- Paperna, I. and R.M. Overstreet. 1981. Parasites and diseases of mullets (Mugilidae). *International Biology Progress* 26:411-493.
- Parker, R.S., C.T. Hackney, and M.F. Vidrine. 1984. Ecology and reproductive strategy of a South Louisiana freshwater mussel, *Glebulula rotundata* (Unionidae: Lampsilini). *Freshwater Invertebrate Biology*. 3(2):53-58.
- Porter, H.J. and K.J. Horn. 1980. Freshwater glochidia from Lake Waccamaw, Columbus County, North Carolina. *Bulletin of the American Malacological Union* 1980:13-17.
- Richard, P.E., T.H. Dietz, and H. Silverman. 1991. Structure of the gill during reproduction in the unionids, *Anodonta grandis*, *Ligumia subrostrata* and *Carunculina parva texasensis*. *Canadian Journal of Zoology* 67:1744-1754.
- Rand, T.G. and M. Wiles. 1982. Species differentiation of the glochidia of *Anodonta cataraacta* Say, 1817 and *Anodonta implicata* Say, 1829 (Mollusca: Unionidae) by scanning electron microscopy. *Canadian Journal of Zoology* 60(7):1722-1727.
- Silverman, H., W.T. Kays, and T.H. Deitz. 1987. Maternal calcium contribution to glochidial shells in freshwater mussels (Eulamellibranchia: Unionidae). *Journal of Experimental Zoology*. 242(2):137-146.
- Smith, D.G. 1976. Notes on the biology of *Margaritifera margaritifera margaritifera* (Lin.) in central Massachusetts. *American Midland Naturalist* 96(1):252-256. 2 figs.
- Smith, D.G. 1978. Biannual gametogenesis in *Margaritifera margaritifera* (L.) in northern North America. *Bulletin of the American Malacological Union* 1978: 49-52.
- Stein, C.B. 1971. Naiad life cycles: their significance in the conservation of the fauna. *Proceedings of a Symposium on Rare and Endangered Mollusks (Naiades) of the U.S.* U.S. Department of the Interior, Bureau of Sport Fisheries and Wildlife. pp. 19-25.
- Stein, C.B. 1973. The life history of *Amblema plicata* (Say, 1817), the three-ridge naiad (Mollusca: Bivalvia). Ph.D. Dissertation, The Ohio State University (Columbus). 216 pp.
- Sterki, V. 1898. *Anodonta imbecilis* hermaphroditic. *The Nautilus* 12:87, 88.
- Surber, T. 1912. Identification of the glochidia of freshwater mussels. *U.S. Bureau of Fisheries Document No.* 771:1-10. Pls. 1-3.
- Surber, T. 1913. Notes on the natural hosts of fresh-water mussels. *Bulletin of the U.S. Bureau of Fisheries* 32:101-116. Pls. 29-31, Tabs. I-VI, 1 ff.
- Surber, T. 1915. Identification of the glochidia of fresh-water mussels. *Bulletin of the U.S. Bureau of Fisheries, Document No.* 813:1-9. Pl. I.
- Tankersley, R.A. and R.V. Dimock, Jr. 1992. Quantitative analysis of the structure and function of the marsupial gills of the freshwater mussel *Anodonta cataraacta*. *Biological Bulletin* 182:145-154.
- Taylor, D.W. and T. Uyeno. 1966. Evolution of host specificity in freshwater salmonid fishes and mussels of the North Pacific region. *Venus, The Japanese Journal of Malacology* 24:199-209.
- Tedla, S. and C.H. Fernando. 1969. Observations on the glochidia of *Lampsilis radiata* (Gmelin) infesting yellow perch, *Perca flavescens* (Mitchell), in the Bay of Quinte, Lake Ontario. *Canadian Journal of Zoology* 47:705-712.
- Tepe, W.C. 1943. Hermaphroditism in *Carunculina parva*, a freshwater mussel. *American Midland Naturalist* 29(3):621-623. 2 figs.
- Tomba, A.S. 1979. Life-cycle completion of the freshwater clam *Lasmigona compressa* (Bivalvia: Unionidae) in an experimental host, *Lebistes reticulatus*. *The Veliger* 22(2):188-190.
- Trdan, R.J. 1981. Reproductive biology of *Lampsilis radiata siliquoidea* (Pelecypoda: Unionidae). *American Midland Naturalist*. 106(2):243-248.
- Trdan, R.J. and W.R. Hoeh. 1982. Eurytopic host use by two congeneric species of freshwater mussel (Pelecypoda: Unionidae: *Anodonta*). *American Midland Naturalist* 108(2):381-388.



- Threlfall, W. 1986. Seasonal occurrence of *Anodonta cataraeta* Say, 1817, glochidia on three-spined sticklebacks, *Gasterosteus aculeatus* Linnaeus. *Veliger* 29(2):231-234.
- Trimble, J.J., II and D. Gaudin. 1975. Fine structure of the sperm of the freshwater clam *Ligumia subrostrata* (Say, 1831) (Mollusca: Bivalvia). *The Veliger* 18(1):34-36. [3 pls.]
- Tucker, M.E. 1927. Morphology of the glochidia and juvenile of the mussel *Anodonta imbecilis*. *Transactions of the American Microscopical Society* 46:286-293. [pl X, tabs. I, II]
- Utterback, W.I. 1916. Parasitism among Missouri naiades. *American Midland Naturalist* 4: 518-521.
- Utterback, W.I. 1933. New glochidia. *Proceedings of the West Virginia Academy of Science* 6:32-36.
- van der Schalie, H. 1970. Hermaphroditism among North American freshwater mussels. *Malacologia* 10(1):93-112.
- van der Schalie, H. and F. Locke. 1941. Hermaphroditism in *Anodonta grandis*, a freshwater mussel. *Occasional Papers of the Museum of Zoology, University of Michigan* No. 432. 7 pp. 3 Pls.
- Waller, D.L., and L.E. Holland-Bartels. 1988. Fish hosts for glochidia of the endangered freshwater mussel *Lampsilis higginsii* Lea (Bivalvia: Unionidae). *Malacological Review* 21(1-2):119-122.
- Waller, D.L., L.E. Holland-Bartels, and L.G. Mitchell. 1988. Morphology of glochidia of *Lampsilis higginsii* (Bivalvia: Unionidae) compared with three related species. *American Malacological Bulletin* 6(1):39-43.
- Waller, D.L., L.E. Holland-Bartels, L.G. Mitchell, and T.W. Kammer. 1985. Artificial infestation of largemouth bass and walleye with glochidia of *Lampsilis ventricosa* (Pelecypoda: Unionidae). *Freshwater Invertebrate Biology* 4(3):152-153.
- Watters, G.T. 1994. An annotated bibliography of the reproduction and propagation of the Unionoidea (Primarily of North America). *Ohio Biological Survey, Miscellaneous Monograph No. 1*. 158 pp.
- Wiles, M. 1974. Parasites of *Fundulus diaphanus* (LeSueur) (Pisces: Cyprinodontidae) in certain Nova Scotian freshwaters. *Canadian Journal of Zoology* 53(11):1578-1580.
- Wiles, M. 1975. The glochidia of certain Unionidae (Mollusca) in Nova Scotia and their fish hosts. *Canadian Journal of Zoology* 53(1):33-41. 5 figs., 11 tabs.
- Wilson, C.B. 1916. Copepod parasites of fresh-water fishes and their economic relations to mussel glochidia. *Bulletin of the Bureau of Fisheries* 34:331-374. [Issued also as Bureau of Fisheries Document No. 824]
- Yeager, B.L. and R.J. Neves. 1986. Reproductive cycle and fish hosts of the rabbit's foot mussel *Quadrula cylindrica strigillata* (Mollusca: Unionidae) in the Upper Tennessee River drainage. *American Midland Naturalist* 116(2):329-340.
- Yeager, B.L. and C.F. Saylor. 1995. Fish hosts for four species of freshwater mussels (Pelecypoda: Unionidae) in the Upper Tennessee River Drainage. *The American Midland Naturalist* 133(1):1-6.
- Yokley, P., Jr. 1972. Life history of *Pleurobema cordatum* (Rafinesque, 1820) (Bivalvia: Unionacea). *Malacologia* 11:351-364.
- Young, D. 1911. The implantation of the glochidium on the fish. *University of Missouri Bulletin Science Series* 2:1-20.
- Zale, A.V. and R.J. Neves. 1982. Fish hosts of four species of Lampsiline mussels (Mollusca: Unionidae) in Big Moccasin Creek, Virginia. *Canadian Journal of Zoology* 60(11):2535-2542.
- Zale, A.V. and R.J. Neves. 1982. Identification of a host fish for *Alasmidonta minor* (Mollusca: Unionidae). *American Midland Naturalist* 107:386-388.
- Zale, A.V. and R.J. Neves. 1982. Reproductive biology of four freshwater mussel species (Mollusca: Unionidae) in Virginia. *Freshwater Invertebrate Biology* 1:17-28.

#### f. Food

- Allen, W.R. 1914. The food and feeding habits of freshwater mussels. I. Experimental studies of the food relations of certain Unionidae. *Biological Bulletin* 27(3):127-147. Plates I-III.
- Allen, W.R. 1921. Studies of the food relations of certain Unionidae. *Biological Bulletin* 40(4):210-241.
- Churchill, E.P., Jr. 1915. The absorption of the fat by fresh-water mussels. *Biological Bulletin* 29(1):68-86, 3 pls.
- Churchill, E.P., Jr. 1916. The absorption of nutrients from solution by fresh-water mussels. *Journal of Experimental Zoology* 21(3):403-429.
- Churchill, E.P., Jr. and S.I. Lewis. 1924. Food and feeding in freshwater mussels. *Bulletin of the U.S. Bureau of Fisheries* 39:439-471 (Figs. 1-26, Tab. 1).
- Coker, R.E., H.W. Clark, A.F. Shira, and A.D. Howard. 1921. Natural history and propagation of freshwater mussels. *Bulletin U.S. Bureau of Fisheries* 37:77-181. Pls. 5-21, Figs. 1-14, Tabs. 1-24.

## APPENDIX 1

Freshwater bivalves reported from the Youghiogheny and Casselman rivers in Pennsylvania and potentially occurring in Maryland<sup>1</sup>.

	Youghiogheny River	Casselman River
<i>Actinonaias ligamentina</i>	HA <sup>2</sup>	-
<i>Amblema plicata</i>	A	-
<i>Cylconaias tuberculata</i>	A	-
<i>Elliptio crassidens</i>	A	-
<i>Elliptio dilatata</i>	A	A
<i>Epioblasma torulosa rangiana</i>	A	-
<i>Fusconaia flava</i>	A	-
<i>Fusconaia subrotunda</i>	A	-
<i>Lampsilis abrupta</i>	A	-
<i>Lampsilis fasciola</i>	A	-
<i>Lampsilis ovata</i>	A	-
<i>Lampsilis siliquioidea</i>	A	-
<i>Lasmigona costata</i>	A	-
<i>Ligumia recta</i>	A	-
<i>Obovaria retusa</i>	A	-
<i>Obovaria subrotunda</i>	A	-
<i>Pleurobema clava</i>	A	-
<i>Pleurobema cordatum</i>	A	-
<i>Pleurobema sintoxia</i>	A	-
<i>Ptychobranhus fasciolaris</i>	A	-
<i>Strophitus undulatus</i>	HA	A
<i>Villosa fabalis</i>	A	-
TOTAL TAXA	22	2

<sup>1</sup> Historic data for the Youghiogheny and Casselman rivers is derived from Ortmann (1919). The archaeological data is from Winters (n.d.), an unpublished manuscript on file in the Div. of Anthropology, Carnegie Museum of Natural History, Pittsburgh, Pennsylvania.

<sup>2</sup> H - Historic record; A - Archaeological record.

